

## Regionalized expression of the *Dbx* family homeobox genes in the embryonic CNS of the mouse

Hiroki Shoji<sup>a,1</sup>, Toshiro Ito<sup>b</sup>, Yoshio Wakamatsu<sup>a</sup>, Naoto Hayasaka<sup>a</sup>, Kanae Ohsaki<sup>c</sup>, Mitsuru Oyanagi<sup>d</sup>, Ryo Kominami<sup>d</sup>, Hisato Kondoh<sup>a</sup>, Naoki Takahashi<sup>c,\*</sup>

<sup>a</sup>Institute for Molecular and Cellular Biology, Osaka University, 1-3, Yamadaoka, Suita, Osaka, 565, Japan

<sup>b</sup>National Institute for Basic Biology, Okazaki, 444, Japan

<sup>c</sup>Graduate School of Biological Sciences, Nara Institute of Science and Technology, 8916-5, Takayama, Ikoma, Nara, 630-01, Japan

<sup>d</sup>First Department of Biochemistry, Niigata University School of Medicine, 1-757, Asahimachi-doori, Niigata, 951, Japan

Received 30 November 1995; accepted 30 January 1996

### Abstract

Here we report the identification of a novel homeobox gene family *Dbx* in mouse, which consists of *Dbx* and *Dbx2*. The two genes share similar structural organization and are encoded by different chromosomes. The predicted *Dbx* and *Dbx2* proteins share 85% identity in their homeodomain amino acid sequences, but otherwise showed no significant similarity. Characterization of the expression of these two genes in the embryos suggested their role in the development of the CNS. In the forebrain, *Dbx* is expressed in various regions, while *Dbx2* showed a more restricted pattern of expression. In the midbrain, the expression domains of *Dbx* and *Dbx2* overlap along the dorso-lateral wall of the ventricle. In the hindbrain and spinal cord, both genes are expressed in the boundary separating the basal and alar plates, which seems to correspond to the sulcus limitans. Expression of the *Dbx/Dbx2* genes is restricted to the ventricular region of the embryonic CNS except for that of *Dbx* in the septum of the telencephalon. Together these observations indicate possible participation of the members of the *Dbx* family in regionalization of the CNS. While the expression of *Dbx* was restricted to the CNS, *Dbx2* was also expressed in some of the mesenchymal cells, such as limb buds and tooth germs.

**Keywords:** Homeobox; Neurogenesis; Dorso-ventral border; Developing brain; Tooth germ

### 1. Introduction

Our understanding of the molecular mechanisms underlying embryonic morphogenesis has been accelerated by genetic studies achieved in *Drosophila melanogaster*. A number of regulatory genes have been identified which have been integrated into a cascade of genetic networks (for review, see Akam, 1987). Homeoboxes were discovered as highly conserved 180 bp sequence motifs shared by a number of genes involved in morphogenesis (McGinnis et al., 1984; Scott and Weiner, 1984). Homeoboxes are translated into homeodomains which serve as DNA binding domains in transcriptional regulation (for

reviews, see Levine and Hoey, 1988; Scott et al., 1989; Hayashi and Scott, 1990; Gehring et al., 1994).

Homeoboxes are highly conserved among various animal species not only in terms of the basic feature of coding amino acid sequences but also in the genomic composition of the homeobox-containing genes. In several animal species, a family of homeobox-containing genes have been identified, and classified into subfamilies based upon the amino acid sequences of their homeodomains. In mouse, nearly a hundred homeobox genes have been identified (Kappen et al., 1993). Murine *Hox* genes are organized into four analogous gene clusters on different chromosomes, and are differentially expressed with individually unique rostro-caudal specificity in various embryonic anlagen depending on the position within the clusters analogous to the *Drosophila Hom-C* (for reviews, see Kessel and Gruss, 1990; Kenyon, 1994; Krumlauf, 1994). Other homeobox genes, such as *Pax* and *Evx1* are

\* Corresponding author.

<sup>1</sup> Present address: Institute for Enzyme Research, University of Tokushima, 3-18-15, Kuramoto, Tokushima, 770, Japan.




Fig. 1. Structure of *Dbx* (A) and *Dbx2* (B) genes. Exon-intron organization maps of genomic clones of *Dbx* (g164) and *Dbx2* (gS4), in comparison with cDNA clones shown below (c164, cS4). Wide boxes indicate the protein coding sequences, open boxes the homeoboxes, and narrow boxes the 3' untranslated regions. a–f indicate probes used for phage library screening (a,c,f), genomic Southern hybridization (b) and in situ hybridization (d,e). BB in the fourth exon of *Dbx* indicates twice repeated *Bam*HI sites separated by a nucleotide between their recognition sequences. Restriction sites: B, *Bam*HI; E, *Eco*RI; H, *Hind*III.

specifically expressed along the dorso-ventral axis of the CNS (Bastian et al., 1990; Gruss and Walther, 1992).

Involvement of the homeobox genes in regional specification of the embryonic anlagen, as suggested by analysis of their embryonic expression patterns, has been demonstrated by the phenotypes of the mutant animals lacking function of one of these genes (for reviews, see Gruss and Walther, 1992; McGinnis and Krumlauf, 1992; Krumlauf, 1994).

The majority of vertebrate homeobox genes have been identified by their similarity to *Drosophila* homeobox genes. In order to identify homeobox genes including those which are highly divergent from *Drosophila* genes, we screened a mouse genomic library using a set of degenerated oligonucleotides corresponding to conserved motif in the homeodomain. This enabled us to isolate a new family of homeobox genes named *Dbx* (developing brain homeobox genes) which consists of *Dbx* (Lu et al., 1992) and *Dbx2*. These two genes are similar in their homeobox sequences and in gene organization, and are both expressed in the embryonic CNS with regional specificities.

## 2. Results

### 2.1. Genomic and cDNA clones of *Dbx* gene family

A  $\lambda$ EMBL3 mouse genomic library was screened by

hybridization with synthetic oligonucleotides representing the most conserved region of the homeoboxes [(A/G)TCAAG(A/G)TCTGGTT(C/T)CA(A/G)AA(C/T)(C/A)GG(C/A)G(C/G)ATGAAGGA] under mild stringency. From  $8 \times 10^5$  plaques, three positive clones, g161, g164 and g169, were obtained. Restriction analysis and nucleotide sequence determination indicated that g161 and g169 corresponded to *HoxB1* (Frohman et al., 1990) and *HoxD3* (Lonai et al., 1987), respectively, while g164 represented a novel gene (Figs. 1A and 2A).

To isolate the corresponding cDNA, we screened a  $\lambda$ gt10 12.5 day mouse embryonic cDNA library using a fragment of g164 as probe. The cDNA clone obtained, c164, was 1795 bp long, contained a poly A tail which was accompanied by a polyadenylation signal (AATAAA) (Figs. 1A and 2A). The longest open reading frame (ORF) found in the c164 sequence began at the 5' end and terminated at nucleotide position 830. Comparison of the c164 and g164 sequences indicated that a putative translational initiation site conforming to Kozak's consensus (Kozak, 1986) existed 171 bp upstream of the 5' end of c164 (Fig. 2A). Translational initiation at this site is consistent with the size of the transcript estimated from Northern analysis (2 kb, data not shown).

To determine whether homeobox sequences related to the g164 gene exist in mouse and chicken genomes, we used a fragment of c164 which included the homeobox (b in Fig. 1) in Southern hybridization analysis. In addition

A

GAATTCAACTATTCTCCGAAACAAAACAAAACAAAACAAACACCCTCAAATCTTCCCAACCTGACGGACTGTCATCGGATGGCACCGG 90  
 AGTTTCTATATTTCCAAGTCTCCCTGGGTGTGGGAGTGGGACTTCTAACCCATCCCTCGCATCTGCTCTACGAGTCCCATTGAGTGAGAA 180  
 ATAGCTGGAGACCGGTATGTTTTCTACTTAAGGGCTGAAATTTCCACACGGCCCTCACACACAAGGGAGAGTTATTTTACCGATAGC 270  
 CTGGAGATGTAGTCTATGAATATTTATTGATTTATTTCTAGTCTCAGCAGGAGCCGCAGATAAATTTAGTGCCTTTAAATAAT 360  
 CCTTTTCTATGAGCAAGATGTCATCTGAACTTCTAAATAGGCATTTAATTAGCCCGGGTGTGCTTCAGAACAAAACCTCCAGCGAG 450  
 AGCTTTGGGGACTCACTGTATAGCAGAGAGCCGTCGATGCACAGCGTTGATGAGCGCCGCTTTGTTGCCGCCCATGTGAGGGCCA 540  
 TGCTTATGAAGACTAATCCAATCATAAATTCATCCCTGAACCAATCAGAGAGCCTGGAGCCTCCCGCAATGAAGCCCGAGTGGAGAAC 630  
 TTTGTTGAACTTCATTGTGAGAGCTGTGAGTTTCAAAGAGCTGGAGCGGGCCACTATAAAACCATCTCCTGGACCGGAGAGCCAC 720  
 AGAAGACTCGCAGACGCCAAGTAGGACGTTACCAGGATACTGAGCCAAAGAGGGCATCGGGGGAGGGGGCGGAAGCCAGGCGCTTGCC 810  
 CTCGCGCCAGGAAAGGGAGCTACTGCCAGACCCGCTCTAACCATGATGTTCCCGGCTCCTTGCGCTCCCGGGGATACCTTAGCC 900  
 M F P G L L A P P A G Y P S L 16

TCCTGCGACCCACGCCACCTTAACACTGCCCCAGTCCCTTCAATCCGCATTTTCCGGCCACTCTAGTTTCTAGTAGAAGATTGATCC 990  
 L R P T P T L T L P Q S L Q S A F S G H S S F L V E D L I R 46

GCATCAGCCGGCTCCACCTACCTGTCCCGCAGCATACCCGCTGCCAGCCTGTACCCCCAGTCAAGAGGCCCTGCAGCCCTCGCCG 1180  
 I S R P P T Y L S R S I P A A S L S P P S Q E A P A A L A D 76

ACTCGGGACCTCAGACCTGGGCTCTCCTGGTCCGGAGCCGGCGAGGAGTTCTCCACAGACCCGCTCTCCCTGCCAGCGAGCCCA 1270  
 S G T S D L G S P G S G S R R G S S P Q T A L S P A S E P T 106

CGTTTCTGAAGTTGGGTGAATGCCATCCTTTCCCTCGCGCTAGAAGAGGTAAGATAACAGCCGTCCTCGACGTGCGTTCCGCGG ( 1358  
 F L K F G V N A I L S S A P R R E 123

-0.5kb-) TCTCTCTCTCTCTCTCTCTCTCTTCCCTCGCCCTACCAGAACTTCGCCCCGTTTGTGTCAGAGCCCTCCTCCCAAGACGTT 1440  
 T S P A L L Q S P P P K T F 137

CGCCTTTCATACTTCGAAGGCTCGTTCCAACCCTTTATCAGATCCCTCCTATTCCCAGGTAGGCTCTTTTCGTGTTTTTCCCGCTCCGG 1530  
 A F P Y F E G S F Q P F I R S S Y F P A 157

GCTGGACAGCCGTTCCAGTCCCGCGGTCGCTCCCTACCCGAAGTCGTATTCTCCGTCCACCCC (-1.5kb-) AAAGGCAATTGT 1610  
 GGCTGCTGTTTCGTGGGTTTTTCTAATCCGGGTTGCACGCTTGAAGCGAAAGTTATTGTGCGGAAAGCGCCTTCACTCAGACAGG 1700  
 TTGTAGGTTCTCCGCTCTATTCTCTACGGGGAGAAAAGGAGTGTTTTAAAGAGATGAAGGGACTCGAGGGGAAAAGGAGGAGAACGC 1790

TAGTAAAACAGCTTGGGAGAGGGGATGCTCGGCTCTGCAGGACACCCGGCTATTCTCTACTTCCCTCACCCCGCAGCTTCTCTCCAGC 1880  
 S S S 160

GTCGTGCTATCCCTGGGACCTTCTCCTGGCCGCTGGCTGCTCGCGCAAGCCCTCGCCGGGCATGCTGCGCAGAGCCGTTCTCTGAC 1970  
 V V P I P G T F S W P L A A R G K P R R G M L R R A V F S D 190

GTGCAGCGGAAAGCCCTGGAGAAGCAGTTCCAGAAGCAGAAGTACATCAGCAAGCCAGACCGGAAAAGCTGGCCCTCAAGCTTGCTTG 2060  
 V Q R K A L E K T F Q K Q K Y I S K P D R K K L A S K L G L 220

AAGGACTCGCAGGTGCGAGCCGCTCCACCG (-0.5kb-) CAAGGCTGGTGGGAAACAACACAGAAATCACCTTGAGGGCAGCGCCTTC 2140  
 K D S Q 224

CTACATGTTCCCAAGCCTGGCTTCTAAGGTTGCCCCAACACCACCACCACCACCACCACCACCACCACCACCACCACCACCACCACC 2230

CCCTCATTGTGAGCAGACACTAACTTGTCTGGTTCTTCCCTCACAGGTGAAAATCTGGTTCCAGAATCGACGCATGAAATGGCGGAAC 2320  
 V K I W F Q N R R M K W R N 238

TCCAAGAGCGGAGCTCCTGTCTAGCGGGGCTGCCAGAGCAGACCCCTTCCACAAAATAAACCCCATCCAGACCTTAGTGATGTA 2410  
 S K E R E L L S S G G C R E Q T L P T K L N P H P D L S D V 268

GGCCAGAAGGGACCTGGGATGAGGAGGAAGACAATCCGGGTGCCCGCTGGCTACCACGCGCCGAGACCCGCGACATTTGCTGGAA 2500  
 G Q K G P G D E E E D N P G A R L A Y H A P A D P R H L L E 298

GGGCCACTGCCTGCTCGCCGGCGACTCGAGCAGCCCGGTAACCGTCAGACTTCTCTGATTCTGATGAGGATGAGGAGGGCGAGGAG 2590  
 G P L P A S P A H S S S P G K P S D F S D S D E D E E G E E 328

GATGAGGAAATCACGGTGTCTAGGAGCCCTCCAGCGGCTCTGGGGCTGTTGCAATTGGAACCTGGAGAGGGAACGCCACGTTGCCAC 2680  
 D E E I T V S \* 335

CTCTGTTCAAGATCCTGGATCCTGCCTTCTCTACAACAGACCCACCACCTTCTCAGGGTGGCATCACCTTCTGTGTGCCAGAGAC 2770  
 ATTTCTTGCCTGCTGCCACCTCTAAACAACCCCTTCCACGGGAGCGTGGGTCTTTTCTACAAGAGCAACATTGCATCTATTTATTG 2860  
 CAAATTTATTTAGACACCGCGTGCAGGGTGTGTGAGAGGGGTGCCCTAGTTCCTACTGCCAGTGTCTACTGTGAGCCAGGTCCTGTCA 2950  
 AACACCAAGCCTTGACCCCTAACTTCTGGATGGCCACCTTCTGAGGGGATCAGTCAGCCATCTGTCTTTACCTTGAGAGTA 3040  
 GAAAGCACAGGAGATTCTGAAAGAGAATAAGGGCACCCTCTCCCTGTCTAAGGGGAAAGCACTATCAAGCAAGCTTTTCAATTCTTGATG 3130  
 CTCCTTTGGCGGGAGGAAGCATTCTCTCTGTAGACATGTGATCCAGTCTGCTTGTATGTCCTTCTGCTACTCCAACGGGAGCCTT 3220  
 TTCAGAGTGCAGCTCCTGGCACTGGGTGTGGACTTGTCTGACTCTCCGACAGTTTCACCCAGCTTTCAGTCTGGGACGCTTACCTG 3310  
 AGATGTGCTCTACTGTATCTATATTTGTATGGATGTTTTATACTGAGGTTATTGTGAATGACCATAATGAAGGGCTGGCTGTGTCC 3400  
 TCTCATTATTTTCTACTGCAATGGTCTTTAAAGTCTAAAAGGTAGCTGTGTAACCTCATAATTTATAAGGAAATGCTTATTTTGTATTTT 3490

TTACCATGTACAGACTTTTTATATATGTATATATGTATCAAATGGACAAATGCCAATAAAAGTAGAATGAGTTTGGCTAAAGGGC 3580  
 ATTTGTAAGTCAAAAACGTGATGGCTGAAAGAACCAGGATTGGTCTTGTGGGTACCCAGAAGGCAA 3647

Fig. 2(A).

B

CACGCACGCCACCCCGCCCGCCACGCCAGCCGCGGGTCTGAGTGTCCGACGCTGCGGCCACTGCCTGCCAGCCAGTCCCGCTCAAGCT 90  
 T H A P P P P R P A P G P E C P Q L R P L P A S P V P L K L 30

GTGCCCGCCGCGGACCCTTTGGTGTGCGCTGGGCCTTCCAGATGCCCCCTGGCCGGGCTCCGGGGGAGCGGGACAGTGCCTTCCAGCCCTC 180  
 C P A G P F G V R W A F Q M P P G R A P G E R D S A F Q P S 60

AGCCCCAG..... [TGCAGGTCAACGGATCATCATTTTTTTGGGTTTCAAGGAAGTCCAGCACTCAGATAGCTTA 248  
 A P G 63

AGCAACAGAGAACAGCCTGCCACCGTGTAACATGAGATACCCTGTCTGAGAGTCCGAGGGCTTCCCCTCTAACCCTGTGCCCTT 338

TTGTATTCTTGTCTCTGCCAGGAGAGAAGAACAGGGCTGCCCTTTGTGTACCAGGACTCAAATCCAAAGCTCGCAGGGGATTTTACGA 428  
 E E H G L P L L T Q D S N S K A R R G I L R 85

AGAGCTGTGTTCTCAGAGGAGCAGAAAAGCTCTGGAGAAAATGTTTTCAGAAAGCAGAAAATACATCAGAAAACAGACCGAAGGAAACTC 518  
 R A V F S E E Q R K A L E K M F Q K Q K Y I S K T D R R K L 115

GCGTTCAGCTTGGGACTGAAGGAGTACAGGTACAGTACAGTACAGTACAGTACAGTACAGTACAGTACAGTACAGTACAGTACAGTACAGT 608  
 A V S L G L K E S Q 125

TTTTGCCAAATATATCTTGTGCCAGGTTTGTACCCCAAAGTGTCTGAAATTTTTCAGTAGGTTGAGGCCCATGGACAAATCCTATGTGAA 698  
 GACAGGATCCACACACGCTGAGGTGGCAAAGCAGACCAGCAATTC (-6.4kb-) CACATTAACACCCCTGCACATAAATGTTGTGTG 778  
 CATATACATACACAGACACCACGTTGTCATCTGCATGTTTTCATGCATGCCACACACACACAAATACATGCACACACACACATACACACAC 868  
ACACACACACACACACACACACACACACACTCCCCGATGCAATCCAATTAATCAGCCAAATCAACACACATAGGTGCTCTACTA 958  
GTCAATTTTCTTGAATAATGTATCAGAATAGAGCAAATATAATGATGTTTAAAAGTATTTAAAATAAGTTAATTTCAACAAGCCACA 1048  
 GAAATACAGAGTGAATCAGTGTGTACACTCATCTTACTTGAAGAAAGTGCCTGTGGTGTGCGGGTTGATTTCTCAAGTCTTTTGAAGGA 1138  
 ACCACGGGAAAAGCCACAGGGAGTCTCTAGTACACTTCCAGAGTAGGAGATTCAGTAAAGTATAGTCAAGCAGTACAGTGGG 1228  
 GATGTTGAAGAGGGATCGGAATATGCAGTTACCCAGCATATATGAGGCCAAAGTCAAAGAAAATAAGGAAAACACTCAAGTGTGTATG 1318

CGGAATAAGGACTTCTTTCTTATATTTTATCATTAGGTTGAAGATTTGGTTCAGAACCAGGATGAAATGGCGGAATCCAAAGAAA 1408  
 V K I W F Q N R R M K W R N S K E K 143

AGGAAGTACTCTCCAGTAGTGTCTCCAAGAAGTAGCCCTTCCAGGAAGACAGGCTGGCAGGCCTGCCGTGGGCTGTCTCCGCAGTGCC 1498  
 E V L S S R C L Q E V S L Q E D R L A R P A V G C P P Q C P 173

CGTCAATATGGGAAGTCTCCAGCCACACTCAAGTCCAAGCTGGAGGGAGGAGACTCCAGAATCTGCAGAAAGACTGACCCAGGAGAATT 1588  
 S I W E V S Q P H S S P S W R E E T P E S A E R L T Q E N S 203

CAGGGTCTGGAAGCAGATTCCTCCAGGTACCTGTATCTGTCTGAGAAAGGACCTAGAGACAAGCATGGACTCCAGAGCACCA 1678  
 G V L E A D S L R G T L Y L C P E K G P R D K H G L Q S T I 233

TTTGACGGAAACATACTCCGTGTACATCTAAAAGAACCCTTAGCCAGTAACACTTGGATTAAAGCCAGTTAGATTGTGCCCTCAAAACTGC 1768  
 \*

CTTAAACTAATACCTTGGCATGATGCCTGAGCTGTGCTTAAGGAGAGCCACTCTACTGTTTCTTATTAGCCCTAGCCCTAAGGCTCAAT 1858  
 GTATATATGGAATAGAACCCTCCACAAAAGTGTGATAAGACTGAAAGAAGAGCCAGGATCAGCCCCAGAGCCTGTTCTGAATATGTGTACA 1948

TATTCGTGTGTCTGTGTGTTGTATGATATGCTCATTAATACATCATCAGAAATTAGTCTCTCCAAGTTTTATATGGAACAAATTCAT 2038  
 TTCATTTTGTATAAGTAAAAAAAAAAAAAAAAACCAACGAGCAAATATCATTTTCTAATAGTGAATCTGGGCTT 2111

Fig. 2. Nucleotide sequences of *Dbx* (A) and *Dbx2* (B) genes. Nucleotides are numbered arbitrary just to indicate the positions in the figure. Predicted amino acid sequences are indicated below the nucleotide sequences with residue numbers shown to the right. The homeodomains are shaded. The (CA)<sub>n</sub> repeat, the regions complementary to the PCR primer set used for chromosomal mapping and the polyadenylation signal are underlined. Arrows indicate the putative splicing sites. Solid arrowhead in (A) indicates the 5' end of c164. Sequences of 5' to [ in (B) are cDNA sequence. Open triangles indicate the poly A addition sites.

to the predicted g164 fragments, several different fragments of mouse DNA hybridized to the probe (data not shown). Multiple hybridizing bands were also obtained with chicken DNA. These results suggested that g164 and related genes form a gene family.

Further screening of the λEMBL3 mouse genomic library with a c164 fragment as probe, resulted in two positive clones. One of them corresponded to *Hlx* (Allen et al., 1991), while the other (clone gS4) represented a novel gene (Figs. 1B and 2B).

<i>Dbx</i>	GMLRRVAVFSDVQRKALEKTFQKQKYISKPDRKKLASKLGLKDSQVKIWFQNRMRKWRNSK	mouse
<i>Dbx2</i>	-I-----EE-----M-----T--R--VS---E-----	85% mouse
<i>CHoxE</i>	-I-----ED-----M-----T-----IN-----E-----	87% chick
<i>h1x-1</i>	-----M-----A-----	97% zebrafish
<i>Hlx</i>	RSWS-----NL---G---R-EI---VT---Q---AM---T-A---V-----H--	68% mouse
<i>H2.0</i>	RSWS-----NL---G---IQ---Q---T---R---AR-N-T-A---V-----HTR	65% fly
<i>AHox1</i>	RKWN-----LM---RG---S---S---VA---E-R---DA-S-T-A-----QEI	63% ascidian
<i>Antp</i>	RKRKRG-QYTRY-TLE---E-HFNRLTRRR-IET-HA-C-TER-I-----KKEN	38% fly

Fig. 3. Amino acid sequences of the homeodomains of *Dbx* and *Dbx2* in comparison with those of *CHoxE* (Rangini et al., 1991), *h1x-1* (Fjose et al., 1994), *Hlx* (Allen et al.), *H2.0* (Barad et al., 1988), *AHox1* (Saiga et al., 1991) and *Antp* (Scott and Weiner, 1984). Similarity score of each homeodomain sequence relative to *Dbx* is shown on the right together with the animal species from which the gene was derived. Dashes indicate the residues identical to *Dbx*.

## Chromosome 15




Fig. 4. Distribution of the haplotypes for 131 progeny from intersub-specific backcross mice. The loci followed in the cross are indicated with arrows to the partial chromosome 15 linkage map. The filled squares represent the C57BL/6 allele, and the open squares represent the MSM allele. Each column represents the chromosome identified in the progeny. The numbers of the progeny carrying each type of chromosome are listed at the bottom. In the linkage map, more proximal markers mapped previously are indicated in smaller type (Dietrich et al., 1994; Suda et al., 1994).

We screened the 12.5 day mouse embryonic cDNA library using a restriction fragment of gS4 as probe. One clone, cS4 consisted of a 1035 bp insert with a 703 bp ORF which was missing an ATG at the 5' end but contained a poly A tail at the 3' end (Figs. 1B and 2B).

The nucleotide sequences and exon-intron boundaries

of the genes represented by g164 and gS4 were reconstructed with the aid of their cDNA sequences, and are shown in Fig. 2. After these sequences were determined, Lu et al. (1992, 1994) reported a new gene *Dbx* (developing brain homeobox gene) which was identical to the gene for g164/c164. There were several sequence disagreements in the 5' and 3' non-coding regions between the cDNA sequence reported by Lu et al. (1994) and g164 (Fig. 2A; nucleotides 799, 3027, 3096–8, 3409, 3415, 3517, 3565–6), which may reflect polymorphism or sequence error. gS4, which was entirely new in the mouse, we named *Dbx2*.

The amino acid sequence conservation between the homeodomains of *Dbx* and *Dbx2* is 85% (Fig. 3). In addition, a significant similarity can be found in the 15 amino acids located C-terminal to the homeodomains. Comparison of the genomic and cDNA clones of *Dbx* indicated that the gene consists of at least four exons. The structure of *Dbx2* has not been completely resolved but the position of the intron/exon junctions in the homeobox was identical to that of *Dbx* (Figs. 1 and 2). The structural similarity and expression specificity to be described below indicated that *Dbx* and *Dbx2* compose a gene family.

Comparison of the amino acid sequences of *Dbx* and *Dbx2* homeodomains with those of other proteins revealed that the zebrafish *hlx-1* (Fjose et al., 1994) and the chicken *CHoxE* (Rangini et al., 1991) appeared to be homologues of *Dbx* and *Dbx2*, respectively. *hlx-1* shared a 97% similarity with *Dbx* and *CHoxE* a 93% similarity with *Dbx2* in the homeodomain (Fig. 3). We have isolated




Fig. 5. Whole-mount in situ hybridization data of 9.5 day embryos probed with *Dbx* (A) and *Dbx2* (B). The procedure for detection of *Dbx* and *Dbx2* was slightly different as described in Experimental Procedures. The stripes of *Dbx* and *Dbx2* expression along the length of rhombencephalon to spinal cord are indicated by arrowheads. In (A), three conspicuous sites of *Dbx* expression are marked: me, mesencephalon; d, a complex of areas from dorsal thalamus to rostral pretectum; ma, mammillary to retromammillary areas. In (B), deposit of the stain in structures other than the CNS was not reproducible. Scale bar, 0.5 mm.




Fig. 6. Expression pattern of *Dbx* (A,C,E) and *Dbx2* (B,D) in 11.5 day embryo in sagittal sections. In situ hybridization data are shown. (A,B), (C,D) are pairs of adjacent sagittal sections. (E) is a medial section parallel to (C). (F) shows hybridization with sense probe. An arrowhead in (D) emphasizes the hybridization signal at the boundary between the dorso-ventral wall and the basal zone of the mesencephalon which may escape from attention. An arrow in (D) indicates the expression of *Dbx2* in the caudal region of the hypothalamus. Lines with labels AB, CD and EF indicate the positions and orientations of transverse and frontal sections in Fig. 7. There may also be a marginal level of expression of *Dbx2* in the cerebral cortex, alar plate of the diencephalon and roof of the telencephalon. Abbreviations: aep, anterior entopeduncular area; bh, basal zone of the hypothalamus; cpt, caudal preteectum; dt, dorsal thalamus; h, hypothalamus; is, rhombencephalic isthmus; m, mesencephalon; ma, mammillary area; oc, optic chiasma; poa, anterior preoptic area; pt, preteectum; rh, rhombencephalon; rpt, rostral preteectum; sc, spinal cord; se, septum; vt, ventral thalamus; zi, zona limitans intrathalamica. Scale bars, 1 mm.

a chicken clone encoding a homeobox sequence that is more homologous to *Dbx* than *Dbx2* or *CHoxE* (data not shown), suggesting that the *Dbx* gene family is conserved evolutionarily in both mouse and chicken genomes.

## 2.2. Chromosomal mapping of *Dbx 2*

The mouse chromosomal location of *Dbx2* was determined by intersubspecific backcross analysis. The 3'




Fig. 7. Frontal (A–D) and transverse (E,F) sections showing expression of *Dbx* (A,C,E) and *Dbx2* (B,D,F) in 11.5 day embryonic brain. (A,B), (C,D), (E,F) are adjacent pairs of sections. Arrows in (B,F) indicates the expression of *Dbx2* in the caudal region of the hypothalamus. The positions and orientations of sections are indicated in Fig. 6. Abbreviations: cge, caudal ganglionic eminence; emt, eminentia thalami; otherwise the same as in Fig. 6. Scale bars, 1 mm. Scales for (E,F) are the same as (A).

intron of the gene contained a (CA)<sub>n</sub> repeat (Fig. 2B) which in many cases gives rise to genetic variation. We synthesized two oligonucleotide primers, 5' CACCCTGC ACATAAACATGTG 3', 5' CTAGTAGAGCACCTATG TGTG 3', that spanned a 164 bp region containing the CA repeat. DNA isolated from C57BL/6 and the MSM strain originating from a Japanese wild mouse, *Mus mus molossinus*, were subjected to polymerase chain reaction (PCR). Analysis of the products by electrophoresis on a polyacrylamide gel exhibited a simple sequence length variation between them. A segregation analysis was then carried out for 131 intersubspecific backcross mice obtained by mating (C57BL/6 × MSM) F<sub>1</sub> to MSM males. The strain distribution pattern was compared to that of microsatellite loci that had been determined by published markers (Dietrich et al., 1994). Fig. 4 summarizes the results of typing backcross mice. A clear linkage of the *Dbx2* locus was found with two markers, D15Mit15 (*HoxC8*) and D15Mit40. From this haplotype analysis, we concluded

that the most likely order of loci on chromosome 15 was cen...D15Mit15-0.8 cM-D15Mit40-6.1 cM-*Dbx2*...telo. Since *Dbx* has been mapped to mouse chromosome 7 (Johnson et al., 1992), *Dbx* and *Dbx2* do not form a gene cluster.

### 2.3. Expression of *Dbx* and *Dbx2* in the embryonic CNS

The spatio-temporal distribution of *Dbx* and *Dbx2* mRNAs was examined by in situ hybridization in embryos from stages older than 6.5 day using *Dbx* or *Dbx2* antisense probes (d and e in Fig. 1). Expression of both genes was detected in the neural tube of embryos of stages later than 9.5 day as shown in embryos hybridized in whole mount (Fig. 5). Remarkably, both *Dbx* and *Dbx2* were expressed in a zone separating alar and basal plates along the length of the spinal cord and rhombencephalon. Expression of *Dbx2* occurred at a lower level and in a broader band than *Dbx* (Fig. 5). *Dbx* expression was also




Fig. 8. Expression pattern of *Dbx2* in the embryonic brain of the stages other than 11.5 day. (A,B) are serial frontal sections of 10.5 day embryonic brain. (C,D) are serial transverse sections of 12.5 day embryonic brain. (E,F) are frontal sections of 13.5 day and 15.5 day embryonic brains, respectively. Arrows in (B,D–F) indicate the expression of *Dbx2* in the caudal region of the hypothalamus. Arrowheads in (E,F) indicates the expression of *Dbx2* in the prepectal area. Signal in the telencephalon close to the bottom of the figure (D) is artifact of the section. An area of shining between the telencephalon and diencephalon in (E) is due to hemocytes rather than to a signal. Abbreviations are the same as in Figs. 6 and 7. Scale bars: (A), 0.5 mm; (C,E), 1 mm. Scale for (F) is the same as (E).

observed in the brain, most conspicuously in the mammillary areas, dorsal thalamus to prepectum, and midbrain.

The regional specificity of *Dbx/Dbx2* expression in the brain was more closely examined in 10.5–15.5 day embryos by in situ hybridization of serial sections through saggital, transverse and frontal planes. Representative examples of in situ hybridization data are shown in Figs. 6–8 and summarized as a scheme in Fig. 12. The pattern of expression of these genes did not substantially change during this period. The expression of *Dbx* and *Dbx2* in the brain was restricted to the ventricular zone except for that of *Dbx* in the septum of the telencephalon to be described below.

### 2.3.1. Expression in the telencephalon

In the telencephalon, *Dbx* expression could be detected in the caudal ganglionic eminence (Fig. 7E) and septum (Fig. 6A), while *Dbx2* was not detectable. *Dbx* expression in the septum was detected in cells out of the ventricular zone.

### 2.3.2. Expression in the diencephalon

In the diencephalon, *Dbx* was expressed mainly in three domains rostral to the dorsal thalamus. One domain covered the anterior entopeduncular area (Fig. 6C,E) and the second domain covered the eminentia thalami (Fig. 7E). The third domain covered the optic chiasma, basal






Fig. 9. Comparison of *Dbx* (C,D) and *Dbx2* (E,F) expression in the spinal cord using adjacent transverse sections of 10.5 day embryos (A,C,E) and 12.5 day embryos (B,D,F). Sections of 10.5 day embryos are at the hind limb level and those of 12.5 day embryos are at a cervical level. (A,B) are phase contrast images of (E,F), respectively. Scale bar, 200  $\mu$ m.

zone of the hypothalamus, mammillary areas (including the retromammillary area) and zona limitans intrathalamica (the boundary separating the dorsal and ventral thalamus) (Figs. 6A,C,E and 7A). Out of these domains, *Dbx*-positive cells scattered among negative cells in the alar zone of the hypothalamus and caudal region of the anterior preoptic area (Figs. 6C, 7A and 12A). In the same diencephalic subdivision *Dbx2* was expressed in a short band located at the alar-basal boundary, from caudal region of hypothalamus to rostral region of the ventral thalamus (arrows in Figs. 6D, 7B,F and 8B,D–F).

Caudal to the zona limitans intrathalamica, *Dbx* expression in the alar zone started near the rostral boundary of the dorsal thalamus and epithalamus, and was extended through the middle of the pretectum (Fig. 6C). In the caudal half of the pretectum, *Dbx* expression was confined to the boundary between alar and basal zones (Fig. 6C). *Dbx2* was expressed in a continuous band along the alar-

basal boundary spanning the dorsal thalamus and the pretectum (Figs. 6D, 7B and 8A,C,E,F). It was noted that expression of both *Dbx* and *Dbx2* increased in the dorsal thalamus away from the zona limitans intrathalamica.

### 2.3.3. Expression in the mesencephalon

In the mesencephalon, *Dbx* was strongly expressed in the dorso-lateral wall, continuously from the pretectum and to the rhombencephalic isthmus (Fig. 6A,C,E). Very weak expression of *Dbx2* could also be seen throughout the dorso-lateral wall overlapping with that of *Dbx* (Figs. 6B,D and 10A). In addition, *Dbx2* was expressed at the boundary between the dorso-lateral wall and the basal zone in its rostral half (arrowhead in Fig. 6D).

In the rhombencephalic isthmus, *Dbx* was expressed at a high level while *Dbx2* was not detected (Fig. 6C,D).




Fig. 10. Expression of *Dbx2* in the limb bud mesenchyme. (A–C) A frontal section of 11.5 day embryo at different magnifications showing expression of *Dbx2* in the hind limb bud. Arrows in (A) indicate the hybridization signal in the dorso-lateral wall of the mesencephalon (m), the hind limb bud (hl) and the spinal cord (sc). (B) is a magnification of the hind limb bud in (A). (D,E) A transverse section of 13.5 day embryonic fore limb bud at level of digits. Arrows in (D) indicate the hybridization signal. I and V in (E) indicate the digit number. (C,E) are phase contrast images of (B,D), respectively. Scale bars: (A), 1 mm; (B), 200  $\mu$ m. Scales for (D,E) are the same as (B).

#### 2.3.4. Expression in the rhombencephalon and spinal cord

Along the length of the neural tube posterior to the rhombencephalic isthmus, the expression of *Dbx* and *Dbx2* was detected as a bilateral pair of narrow bands at the boundary between the basal and alar plates (Figs. 5, 6, 7C–F, 8C and 9).

In the spinal cord, expression of *Dbx* was restricted to the boundary between the alar and basal plates (Figs. 5A and 9C,D). Expression of *Dbx2* was similar to *Dbx* in its distribution, but broader, extending out from both sides of the *Dbx*-expressing band and with less defined borders (Figs. 5B and 9E,F). In the rhombencephalon, expression of both genes was restricted to the boundary of the dorso-ventral zones in a manner similar to that seen in the spinal cord (Fig. 5, 7C–F, and 8C).

At early stages when lateral plates of the spinal cord

still consisted of neuroepithelial germinal cells (9.5–10.5 day), expression of *Dbx* and *Dbx2* can be detected throughout the entire thickness of the plates (Fig. 9C,E). At later stages when neural differentiation initiates in the intermediate zone, expression of both genes becomes confined to the ventricular zone (Fig. 9D,F). But regardless of the developmental stage, the dorso-ventral width of the *Dbx/Dbx2*-expressing bands remained fairly the same. Expression of *Dbx2* appears to increase with the development of the embryo as judged by the increase in intensity of the hybridization signal (Fig. 9E,F).

#### 2.4. Expression of *Dbx 2* outside of the CNS

Although expression of *Dbx* was confined to the CNS, *Dbx2* was also expressed in mesenchymal components of the limb buds and tooth germs.




Fig. 11. Expression of *Dbx2* in the tooth germ mesenchyme. (A–C) and (D–F) are frontal sections of 13.5 day and 15.5 day embryonic head, respectively. (B,E) are magnifications of the tooth germs indicated with arrowheads in (A,D), respectively. Arrows in (A,D) show expression of *Dbx2* in the other tooth germs. (C,F) are phase contrast images of (B,E), respectively. The region between two arrowheads in (F) is dental ledge. Signal in lens seen in (A) is non-specific. Abbreviations: df, dental follicle; dp, dental papilla; ie, internal enamel epithelia. Scale bars: (A), 1 mm; (B), 200  $\mu$ m.

#### 2.4.1. Expression in the limb buds

In the limb bud mesenchyme, a significant level of *Dbx2* expression occurred after 11.5 day (Fig. 10A–C). High expression appears localized to the internal mesenchymal cells which later participate in chondrogenesis. After 13.5 day the expression was localized to the cells surrounding the precartilaginous condensation of the digits (Fig. 10D,E).

#### 2.4.2. Expression in the tooth germs





In the tooth germs, *Dbx2* was expressed in the mesenchymal cells surrounding the enamel organ. Data of 13.5 and 15.5 day embryos are shown in Fig. 11. The *Dbx2*-positive cells included both dental papilla and follicle cells, but not the cells immediately adjacent to the internal enamel epithelia of the enamel organ and the cells surrounding the dental ledge. The expression pattern did not substantially change during this period.

### 3. Discussion

#### 3.1. *Dbx* gene family

We have described the gene organization and expression of two new mouse homeobox genes which compose the *Dbx* family. *Dbx* and *Dbx2* have similar homeodomain sequences and exon-intron organizations. *Dbx* cDNA was cloned independently by Lu et al. (1992) and its gene organization reported (Lu et al. 1994), which are in general agreement with our results.

The homeodomain and C-proximal flanking amino acid sequences are very similar between *Dbx* and *Dbx2*, but the sequences of other regions are quite divergent. *Dbx* and *Dbx2* likely can recognize analogous nucleotide sequences, but may differ in their action as transcriptional regulators. Proximal to the C terminus of *Dbx* protein, there is a cluster of acidic residues often present in tran-



scriptional activators (Sigler, 1988), but this motif is absent in *Dbx2*.

Comparison of the homeodomain amino acid sequences indicated that the *Dbx* family is related, but apparently distinct from the *H2.0* family (Fig. 3). Similarity of 63–68% was found between the *Dbx* homeodomain and the homeodomains of the *H2.0* family: *H2.0* (Barad et al., 1988), *Hlx* (Allen et al., 1991) and *AHox1* (Saiga et al., 1991). Invertebrate members for the *Dbx* family have not yet been reported.

### 3.2. Expression of *Dbx* and *Dbx2* in the embryonic brain

*Dbx* and *Dbx2* are expressed in the embryonic brain with regional specificities as summarized in Fig. 12A,B. Expression of *Dbx* is mapped in various regions of the diencephalon with boundaries largely consistent with the neuromeric model of the embryonic brain of the mouse proposed by Bulfone et al. (1993), by Puelles and Rubenstein (1993) and by Rubenstein et al. (1994). *Dbx* expression in the brain had also been reported by Lu et al. (1992, 1994), but not in great detail. *Dbx2* expression in the diencephalon is limited to the alar-basal boundaries of the caudal hypothalamus to the region rostral to the ventral thalamus, dorsal thalamus and pretectum. In the mid-brain, expression of *Dbx* and *Dbx2* largely overlaps along the dorso-lateral wall.

It should be noted that expression of the *Dbx* genes emphasizes the separation of the alar and basal compartments. It is also remarkable that *Dbx* expression in the alar compartment of the pretectum continuous from the dorsal thalamus terminates in the middle of the pretectum with clear demarcation. This indicates that pretectum may be divided into rostral and caudal components as proposed by Figdor and Stern (1993). Thus, the *Dbx* genes provide two important genetic landmarks to the generic map of the brain.

### 3.3. Expression in the spinal cord

Along the length of the neural tube posterior to the rhombencephalic isthmus, both *Dbx* and *Dbx2* are expressed at the boundary of the alar and basal plates which later develop to sensory and motor columns, respectively (Fig. 12C,D). It is important to note that these genes are expressed only in mitotically active cells of the ventricular zone in these regions and that these genes encode pu-

tative transcriptional regulators. It is possible that the *Dbx* family plays a role in specifying a small population of neural cells at the boundary between the columns, a region classically called the sulcus limitans. Expression of these genes overlaps, but the expression of *Dbx2* is broader in a dorso-ventral direction. Therefore, the cells expressing both *Dbx* and *Dbx2* are flanked by cells expressing only *Dbx2*. The overlapping expression of the genes may indicate functional redundancy, but we prefer the view that the products of these genes have different transcriptional regulational activities, as suggested by the high divergence outside of the homeodomain.

It is interesting to speculate a possible function for the sulcus limitans cells in relation to expression of *Dbx* genes. It is known that the differentiation of motor neurons depends on inducing signals from the notochord and floor plate (Placzek et al., 1991; Yamada et al., 1991). Cells expressing *Dbx* genes may acquire a capacity to halt the signal from the ventral region which otherwise induces motor neurons throughout the spinal cord. Altman and Bayer (1984) have proposed that the region of the sulcus limitans (called the intermediate plate) is a source of contra- and ipsilaterally projecting relay neurons. The *Dbx* family may contribute to the determination of cell fates of these neurons. In either case, it is interesting to note that a certain class of cadherins is expressed in ventricular zone cells of the sulcus limitans similar to the *Dbx* family (M. Takeichi, personal communication). It is an intriguing possibility that the *Dbx* family of genes may regulate cadherin expression, which modifies cell interactions and consequently alters fate of the neuroepithelial cells.

### 3.4. Expression of homologues in other animal species

Zebrafish *hlx-1*, a putative homologue of *Dbx*, shows an expression pattern similar to *Dbx* in the CNS (Fjose et al., 1994). However, prior to expression in the CNS, *hlx-1* is expressed in an area of primitive mesoderm (Fjose et al., 1994) which was not observed in the mouse embryo.

Chicken *CHoxE*, the putative chicken homologue of *Dbx2*, has been reported to be expressed in the developing neural tube posterior to the rhombencephalic isthmus almost identically to *Dbx2*. But there is no description of more anterior expression or expression in the other organs (Rangini et al., 1991).

---

Fig. 12. Schematic illustrations of *Dbx* and *Dbx2* expression patterns. (A,B) Expression of *Dbx* and *Dbx2*, respectively, in the brain of 11.5–12.5 day embryo. Gray areas indicate expression domains. The intensity of these paints reflect the relative intensity of hybridization signals in each illustration. Shaded area indicates scattering of *Dbx*-positive cells. (C,D) The spinal cord of 9.5–10.5 day embryos (C) and 11.5–12.5 day embryo (D). Filled areas are the regions expressing both *Dbx* and *Dbx2*, and shaded areas the regions expressing only *Dbx2*. Abbreviations: aep, anterior entopeduncular area; ap, alar plate; bh, basal zone of the hypothalamus; bp, basal plate; cge, caudal ganglionic eminence; cpt, caudal pretectum; dt, dorsal thalamus; emt, eminentia thalami; et, epithalamus; fp, floor plate; h, hypothalamus; is, rhombencephalic isthmus; lge, lateral ganglionic eminence; m, mesencephalon; ma, mammillary area; mge, medial ganglionic eminence; oc, optic chiasma; os, optic stalk; poa, anterior preoptic area; r1–r8, rhombomere 1–8; rm, retromammillary area; rp, roof plate; rpt, rostral pretectum; sc, spinal cord; se, septum; vt, ventral thalamus; vz, ventricular zone; zi, zona limitans intrathalamica.

### 3.5. Complementary expression territories with other homeobox genes

The proposed neuromere map of the CNS (Bulfone et al., 1993; Puelles and Rubenstein, 1993; Rubenstein et al., 1994) is based on the expression patterns of various homeobox genes. Examination of the expression of such genes indicated that some are expressed in regions of brain complementary to *Dbx/Dbx2* expression. For instance, *Pax-6* is expressed roughly complementary to *Dbx* in the diencephalon (Walther and Gruss, 1991; Puelles and Rubenstein, 1993). Caudal to rhombomere 6 of the rhombencephalon and spinal cord, *Gbx-2* is expressed both in alar and basal plates, but not in the sulcus limitans (Bulfone et al., 1993), showing a complementarity to *Dbx* family expression. These observations may suggest that characteristics of individual neuromeres are defined by multiple superpositions of dichotomous choice of gene expression. It is interesting to see if complementary pairs of transcription factor genes are generally found with boundaries which conform to the model of the CNS.

### 3.6. Expression in the limb buds and teeth

Both in the limb buds and tooth germs, *Dbx2* is expressed in undifferentiated mesenchymal cells. In the limb buds, *Dbx2* seems to be expressed in the cells prior to participating in the precartilaginous condensation. In the tooth germs, *Dbx2* is expressed in the mesenchymal cells surrounding the enamel organ, but not in the cells immediately adjacent to the internal enamel epithelia which may be undergoing differentiation into odontoblasts. Therefore, in the mesenchymal cells of these organs, *Dbx2* seems to be expressed transiently immediately before the start of differentiation, which may suggest that *Dbx2* is playing a role in determination of these cells. This is also consistent with the observation in the CNS that *Dbx2* is exclusively expressed in the undifferentiated cells of the ventricular zone.

## 4. Experimental procedures

### 4.1. Genomic and cDNA libraries

BALB/c mouse liver DNA was partially digested with *Sau3AI* and fractionated by sucrose density gradient centrifugation. Fragments with the size of 15–20 kb were ligated to  $\lambda$ EMBL3 vector arms (Clontech) and packaged in vitro (Stratagene). cDNAs for polyA<sup>+</sup> RNAs from 12.5 day ICR mouse embryos were methylated by *EcoRI* methylase, ligated to *EcoRI* linkers and digested with *EcoRI*. cDNAs larger than 500 bp were selected by Sepharose-CL4B column chromatography, and cloned into a  $\lambda$ gt10 vector. Screening of the genomic library using the synthetic oligonucleotide probe was done in hybridization buffer of 6  $\times$  SSC, 0.1% SDS, 0.2% Denhardt's solution,

100  $\mu$ g/ml denatured salmon sperm DNA, at 55°C overnight, followed by three washes with 2  $\times$  SSC, 0.1% SDS for 15 min at 55°C. Screening for isolation of *Dbx2* genomic clone (gS4) was done using as probe the 0.56 kb fragment of *Dbx* cDNA clone (c164) (c in Fig. 1) in the same buffer at 65°C, followed by two washes with 1  $\times$  SSC, 0.1% SDS for 30 min at 65°C. Screening of the cDNA library using a 0.75 kb *EcoRI-BamHI* fragment of g164 or a 0.2 kb *EcoRI* fragment of gS4 as probes (a and f in Fig. 1) was done in the same hybridization condition as described above for gS4, followed by two washes with 0.1  $\times$  SSC, 0.1% SDS for 30 min at 65°C.

### 4.2. DNA sequencing

DNA was sequenced by the dideoxy method (Sanger et al., 1977) using Sequenase Version 2.0 DNA Sequencing Kit (US Biochemical).

### 4.3. Chromosomal localization

Intersubspecific backcross progeny were generated by mating (C57BL/6  $\times$  MSM) F<sub>1</sub> females to MSM males (kindly provided by Drs. K. Moriwaki and N. Miyashita). DNA isolation, PCR and polyacrylamide gel electrophoresis were performed as described (Dietrich et al., 1994). The primer sequences used for D15Mit15 and D15Mit40 were the same as described (Dietrich et al., 1994).

### 4.4. In situ hybridization

*In situ* hybridization with sections was done essentially as described previously (Wakamatsu and Kondoh, 1990). RNA probes were synthesized in vitro by transcription of a linear DNA template of 3' untranslated regions of c164 and gS4 (d and e in Fig. 1) using 200  $\mu$ Ci [<sup>35</sup>S]UTP and T7 RNA polymerase (Toyobo). Paraffin sections of 6.5–15.5 day embryos of ICR mice were prepared on glass slides. The slides were treated with 0.2 N HCl and proteinase K, and then acetylated. Hybridization was done for 20 h at 50°C in 50% formamide, 0.2% Denhardt's solution, 2  $\times$  SSC, 10 mM EDTA, 10% Dextran sulfate, 500  $\mu$ g/ml denatured salmon sperm DNA, 500  $\mu$ g/ml yeast tRNA, 10 mM DTT, and with <sup>35</sup>S-labelled RNA probes. After hybridization, the slides were treated with 20  $\mu$ g/ml RNase A and 1 U/ml RNase T1 at 37°C for 30 min, and washed in 0.1  $\times$  SSC, 20% formamide, 10 mM DTT at 50°C overnight. Autoradiography was done by exposing the slides to Kodak NTB-2 emulsion for 1–2 weeks at 4°C and by developing with Konicadol X (Konica).

Whole-mount *in situ* hybridization of 9.5 day embryos was performed as described by Sasaki and Hogan (1993) for *Dbx*. However, hybridization signal with *Dbx2* was too faint to detect, so a procedure described by Kobayashi et al. (1994) was employed to improve signal detection. A

digoxigenin labeled uridine for RNA probe was purchased from Boehringer Mannheim.

### Acknowledgements

We thank Professor H. Fujisawa for his guidance in neuroanatomy of the embryos, Professor M. Takeichi for information of cadherin expression, Professor S. Noji for advice in whole-mount in situ hybridization and Dr. Ruth Yu for critical reading. This work was supported by grants from the Ministry of Education, Science and Culture of Japan and from Senri Life Science Foundation to H. Kondoh and N. Takahashi.

### References

- Akam, M. (1987) *Development* 101, 1–22.
- Allen, J.D., Lints, T., Jenkins, N.A., Copeland, N.G., Strasser, A., Harvey, R.P. and Adams, J.M. (1991) *Genes Dev.* 5, 509–520.
- Altman, J. and Bayer, S.A. (1984) In Beck, F., Hild, W., van Limborgh, J., Ortman, R., Pauly, J.E. and Schiebler, T.H. (Eds), *Advances in Anatomy, Embryology and Cell Biology*, Vol. 85, Springer Verlag, Berlin.
- Barad, M., Jack, T., Chadwick, R. and McGinnis, W. (1988) *EMBO J.* 7, 2151–2161.
- Bastian, H. and Gruss, P. (1990) *EMBO J.* 9, 1839–1852.
- Bulfone, A., Puelles, L., Porteus, M.H., Frohman, M.A., Martin, G.R. and Rubenstein, J.A.R. (1993) *J Neurosci.* 13, 3155–3172.
- Dietrich, W.F., Miller, J.C., Steen, R.G., Merchant, M., Damron, D., Nahf, R., Gross, A., Joyce, D.C., Wessel, M., Dredge, R.D., Marquis, A., Stein, L.D., Goodman, N., Page, D.C. and Lander, E.S. (1994) *Nature Genet.* 7, 220–225.
- Figdor, M.C. and Stern, C.D. (1993) *Nature* 363, 630–634.
- Fjose, A., Izpisua-Belmonte, J.C., Fromental-Ramain, C. and Duboule, D. (1994) *Development* 120, 71–81.
- Frohman, M., Boyle, M. and Martin, G.R. (1990) *Development* 110, 589–607.
- Gehring, W.G., Qian, Y.Q., Billeter, M., Furukubo-Tokunaga, K., Schier, A.F., Resendez-Perez, D., Affolter, M., Otting, G. and Wüthrich, K. (1994) *Cell* 78, 211–223.
- Gruss, P. and Walther, C. (1992) *Cell* 69, 719–722.
- Hayashi, S. and Scott, M.P. (1990) *Cell* 63, 883–894.
- Johnson, K.R., Lu, S., Murtha, M.T., Ruddle, F.H. and Davisson, M.T. (1992) *Genomics* 14, 1107–1109.
- Kappen, C., Schughary, K. and Ruddle, F.H. (1993) *Genomics* 18, 54–70.
- Kenyon, C. (1994) *Cell* 78, 175–180.
- Kessel, M. and Gruss, P. (1990) *Science* 249, 374–379.
- Kobayashi, S., Saito, H. and Okada, M. (1994) *Dev. Growth Differ.* 36, 629–632.
- Kozak, M. (1987) *Nucleic Acids Res.* 15, 8125–8146.
- Krumlauf, R. (1994) *Cell* 78, 191–201.
- Levine, M. and Hoey, T. (1988) *Cell* 55, 537–540.
- Lonai, P., Arman, E., Czosnek, H., Ruddle, F.H. and Blatt, C. (1987) *DNA* 5, 409–418.
- Lu, S., Bogarad, L.D., Murtha, M.T. and Ruddle, F.H. (1992) *Proc. Natl. Acad. Sci. USA* 89, 8053–8057.
- Lu, S., Wise, T.L. and Ruddle, F.H. (1994) *Mech. Dev.* 47, 187–195.
- McGinnis, W. and Krumlauf, R. (1992) *Cell* 68, 283–302.
- McGinnis, W., Levine, M.S., Hafen, E., Kuroiwa, A. and Gehring, W.J. (1984) *Nature* 308, 428–433.
- Placzek, M., Yamada, T., Tessier-Lavingne, M., Jessell, T.M. and Dodd, J. (1991) *Development Suppl.* 2, 105–122.
- Puelles, L. and Rubenstein, L.R. (1993) *TINS* 16, 472–479.
- Rangini, Z., Ben-Yehuda, A., Shapira, E., Gruenbaum, Y. and Fainsod, A. (1991) *Mech. Dev.* 35, 13–24.
- Rubenstein, J.L.R., Martinez, S., Shimamura, K. and Puelles, L. (1994) *Science* 266, 578–580.
- Saiga, H., Mizokami, A., Makabe, K.W., Satoh, N. and Mita, T. (1991) *Development* 111, 821–828.
- Sanger, F., Nicklen, S. and Coulson, A.R. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463–5467.
- Sasaki, H. and Hogan, B.L.M. (1993) *Development* 118, 47–59.
- Scott, M.P. and Weiner, A. (1984) *Proc. Natl. Acad. Sci. USA* 81, 4115–4119.
- Scott, M.P., Tamkun, J.W. and Hartzell, III, G.W. (1989) *Biochem. Biophys. Acta* 989, 25–48.
- Sigler, P.B. (1988) *Nature* 333, 210–212.
- Suda, T., Oyanagi, M., Wakana, S., Takahashi, Y., Kanada, H., Yonekawa, H., Miyashita, N., Shiroishi, T., Moriwaki, K. and Kominami, R. (1994) *DNA Res.* 1, 169–174.
- Wakamatsu, Y. and Kondoh, H. (1990) *Acta Histochem. Cytochem.* 23, 367–374.
- Walther, C. and Gruss, P. (1991) *Development* 113, 1435–1449.
- Yamada, T., Placzek, M., Tanaka, H., Dodd, J. and Jessell, T.M. (1991) *Cell* 64, 635–647.