

A questionnaire survey on radiation protection among 282 medical staff from 26 endoscopy-fluoroscopy departments in Japan

Shiro Hayashi^{1,2}  | Mamoru Takenaka³ | Hirofumi Kogure⁴ | Takayuki Yakushijin⁵ | Hirotugu Maruyama⁶  | Yasuki Hori⁷ | Toshiyuki Yoshio⁸  | Kenji Ikezawa⁹ | Tadayuki Takagi¹⁰ | Satoshi Asai¹¹ | Kazuhiro Matsunaga¹² | Kengo Matsumoto² | Hidetaka Tsumura¹³ | Shinjiro Yamaguchi¹⁴ | Tetsuya Sumiyoshi¹⁵  | Koji Nagaike¹⁶ | Yuzuru Tamaru¹⁷ | Kazuo Hara¹⁸ | Toshio Fujisawa¹⁹ | Ichiro Oda²⁰ | Ken Ohnita²¹ | Motohiko Kato²² | Hiroko Nebiki²³ | Tatsuya Mikami²⁴ | Akihiro Nishihara²⁵ | Satoshi Egawa²⁶ | Ryuki Minami²⁷ | Makoto Hosono²⁸ | Tsutomu Nishida² 

¹ Department of Gastroenterology and Internal Medicine, Hayashi Clinic, Suita, Osaka, Japan

² Department of Gastroenterology, Toyonaka Municipal Hospital, Toyonaka, Osaka, Japan

³ Department of Gastroenterology and Hepatology, Kindai University Faculty of Medicine, Osakasayama, Osaka, Japan

⁴ Department of Gastroenterology, The University of Tokyo, Tokyo, Japan

⁵ Department of Gastroenterology and Hepatology, Osaka General Medical Center, Osaka, Osaka, Japan

⁶ Department of Gastroenterology, Osaka City University Graduate School of Medicine, Osaka, Osaka, Japan

⁷ Department of Gastroenterology and Metabolism, Nagoya City University Hospital, Nagoya, Aichi, Japan

⁸ Department of Gastroenterology, Cancer Institute Hospital, Tokyo, Japan

⁹ Department of Hepatobiliary and Pancreatic Oncology, Osaka International Cancer Institute, Osaka, Osaka, Japan

¹⁰ Department of Gastroenterology, Fukushima Medical University School of Medicine, Fukushima, Japan

¹¹ Department of Gastroenterology, Tane General Hospital, Osaka, Japan

¹² Department of Gastroenterology, Ishikawa Prefectural Central Hospital, Kanazawa, Ishikawa, Japan

¹³ Department of Gastroenterological Oncology, Hyogo Cancer Center, Akashi, Hyogo, Japan

¹⁴ Department of Gastroenterology and Hepatology, Kansai Rosai Hospital, Amagasaki, Hyogo, Japan

¹⁵ Department of Gastroenterology, Tonan Hospital, Sapporo, Hokkaido, Japan

¹⁶ Department of Gastroenterology and Hepatology, Suita Municipal Hospital, Suita, Osaka, Japan

¹⁷ Department of Gastroenterology, Kure Medical Center and Chugoku Cancer Center, Kure, Hiroshima, Japan

¹⁸ Department of Gastroenterology, Aichi Cancer Center, Nagoya, Aichi, Japan

¹⁹ Department of Gastroenterology, Juntendo University, Bunkyo-ku, Tokyo, Japan

²⁰ Endoscopy Division, National Cancer Center Hospital, Tokyo, Japan

²¹ Department of Gastroenterology and Hepatology, Shunkaikai Inoue Hospital, Nagasaki, Japan

²² Department of Gastroenterology, Keio University School of Medicine, Shinjuku-ku, Tokyo, Japan

²³ Department of Gastroenterology, Osaka City General Hospital, Osaka, Osaka, Japan

²⁴ Division of Endoscopy, Hirosaki University Hospital, Hirosaki, Aomori, Japan

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *DEN Open* published by John Wiley & Sons Australia, Ltd on behalf of Japan Gastroenterological Endoscopy Society

²⁵ Department of Gastroenterology, Minoh City Hospital, Mino, Osaka, Japan

²⁶ Department of Gastroenterology, Osaka Police Hospital, Osaka, Japan

²⁷ Department of Gastroenterology, Tenri Hospital, Tenri, Nara, Japan

²⁸ Department of Radiology, Kindai University Faculty of Medicine, Osakasayama, Osaka, Japan

Correspondence

Tsutomu Nishida, MD PhD, Department of Gastroenterology, Toyonaka Municipal Hospital, 4-14-1 Shibahara, Toyonaka, Osaka 560-8565 Japan.

Email: tnishida.gastro@gmail.com

Abstract

Background and aims: It is essential for endoscopists, technologists, and nurses to understand radiation protection. However, protective equipment usage is still low, and there is little awareness of radiation protection in practice.

Methods: We conducted a questionnaire survey on radiation protection from January to February 2020. The participants were medical staff, including medical doctors, nurses, and radiological and endoscopy technician in endoscopy-fluoroscopy departments. The questionnaire included 14 multiple-choice questions divided among three parts: background, equipment, and knowledge.

Results: We surveyed a total of 282 subjects from 26 institutions. There were 168 medical doctors (60%), 90 nurses (32%), and 24 technologists (9%). Although almost all staff members (99%) always wore a lead apron, only a few wore a thyroid collar (32%) and lead glasses (21%). The rate of wearing a radiation dosimeter was insufficient (69%), especially among doctors (52%). A few subjects knew the radiation exposure dose of each procedure (15%), and slightly over half had attended lectures on radiation protection (64%) and knew about the three principles of radiation protection (59%). Protection adherence did not differ by years of experience, knowledge of fluoroscopy, awareness of radiation exposure doses, or attendance at basic lectures on radiation protection. However, medical doctors who were aware of the radiation exposure dose of each procedure were significantly more likely to wear dosimeters than those who were not ($p = 0.0008$).

Conclusion: Medical staff in endoscopy departments in Japan do not have enough radiation protection equipment or education.

KEYWORDS

education, endoscopy, medical staff, questionnaire survey, radiation protection

INTRODUCTION

Radiation protection is the basis for the safety of both patients and medical staff due to its adverse effects represented by carcinogenicity and skin disorder.^{1,2} The International Commission on Radiological Protection (ICRP) stated that an understanding and awareness of the hazards of radiation among medical staff can prevent unnecessary risks for the population as a whole.^{3,4} In the field of gastroenterology, the World Gastroenterology Organization practice guidelines and the European Society of Gastroenterology Endoscopy guidelines state the importance of radiation protection.^{5,6} However, some reports from Ireland, Korea, and the United States still showed low protective equipment usage and little awareness of radiation protection in practice.⁷⁻⁹ The use of protective equipment and awareness of radiation protection do not

appear to be widespread. Currently, we are conducting a prospective multicenter study in Japan (REX-GI study, UMIN000036525) that will involve the collection of actual radiation exposure-related data from digestive endoscopy interventional procedures.¹⁰ To complement this study, we conducted a nationwide questionnaire survey to assess the actual rate of equipment usage and knowledge and awareness of radiation protection among staff at institutions related to the REX-GI study group and the Fight Japan study group, including gastroenterology doctors, nurses, and technologists from endoscopy-fluoroscopy departments in Japan.

MATERIALS AND METHOD

We conducted a questionnaire survey on radiation exposure protection and collected responses from January

TABLE 1 Questionnaire questions and answers (participants' responses were anonymous)

Question	Answer
1. What is your gender?	a) Female, b) Male
2. How old are you?	a) Twenties, b) Thirties, c) Forties, d) Fifties, e) Over sixty
3. What is your job title?	a) Medical doctor, b) Nurse, c) Technologist
4. What is the size of your institution?	a) University hospital or medical center, b) Regional general hospital (>300 beds), c) Other
5. How many years of career experience do you have?	a) 1–5, b) 6–10, c) 11–15, d) 16–20, e) Over 21 years
6. Do you operate the fluoroscopy unit?	a) Yes, b) No
7. Do you always wear a lead apron?	a) Yes, b) No
8. Do you always wear a thyroid collar?	a) Yes, b) No
9. Do you always wear lead glasses?	a) Yes, b) No
10. Do you always wear a radiation dosimeter?	a) Yes, b) No
11. What type is your fluoroscopy unit, an under-couch or over-couch C-arm system?	a) Under-couch (exposure from below), b) Over-couch (expose from above), c) I don't know
12. Do you know how much radiation dose you are exposed to in each endoscopic procedure under fluoroscopy?	a) Yes, b) No
13. Have you ever attended a basic lecture on radiation exposure?	a) Yes, b) No
14. Do you know the three principles of radiation protection?	a) Yes, b) No

2020 to February 2020. We emailed a representative doctor in each institution involved in the REX-GI study¹⁰ and Fight Japan study group, as well as at other institutions in Japan and invited them to participate in the questionnaire survey. If the representative accepted our invitation, he or she also asked the medical staff, doctors, nurses, and radiological and endoscopy technicians from the fluoroscopic endoscopy suites at each institution to answer an anonymous online questionnaire using Google Forms. Endoscopy technician is a job to maintain the endoscopic equipment and to support the endoscopic procedures in the endoscopy unit. Participants provided informed consent by opening the survey.

The questionnaire used in the survey included 14 multiple-choice questions divided among the following three parts: background, equipment, and knowledge. The details of the questionnaire details are shown in Table 1. Briefly, questions 1–6 regard the background of each person or institution. Questions 7–10 asked about the proper equipment for radiation protection. Questions 11–14 focused on knowledge of radiation exposure and protection. We counted the number of responses from subjects with each job title and compared the numbers of responses among job titles. We then investigated how years of career experience, facility size, and knowledge influenced protective behaviors among doctors.

Statistical analysis

The categorical variables are expressed as the number in each category or the frequency and were compared using the chi-square test or Fisher's exact test when appropriate. A p-value of 0.05 was considered to indi-

cate statistical significance. All statistical analyses were performed with JMP software (ver. 14.3, SAS Institute Inc., Cary, NC, USA).

RESULTS

Responses to the questionnaire

We emailed survey invitations to one hundred and eleven institutions. We obtained answers from two hundred eighty-two subjects, including endoscopists, nurses, and technicians at 26 institutions (participation rate: 26/111, 23.4%).

Questions 1–6: There were 166 (59%) males. Most of the subjects were in their 30s (105, 37%) or 40s (103, 37%). Doctor was the most common occupation (168, 60%). A total of 135 subjects worked at university hospitals or medical center hospitals (48%), 110 worked at regional general hospitals (>300 beds) (39%), and the other 37 worked at other types of hospitals (13%). Regarding years of experience, 45 (16%) had 1–5 years and 68 (24%), 56 (20%), and 56 (20%) had 6–10, 15–20, and over 21 years, respectively. One hundred eighty-eight subjects (67%) had operated fluoroscopy units by themselves (Table 2). Questions 7–10: Two hundred eighty-one subjects (99%) always wore a lead apron, 90 subjects (32%) wore a thyroid collar, 59 subjects (21%) wore lead glasses, and 194 subjects (69%) wore a radiation dosimeter. Questions 11–14: Thirty subjects (11%) did not know the type of fluoroscopy unit. Forty-two subjects (15%) were aware of the radiation dose of each procedure, 180 subjects (64%) had received lectures on radiation protection, and 167 subjects (59%)

TABLE 2 Answers for all subjects including medical doctors, nurses, and technologists

Questions	Answer	All N = 282	Medical doctors N = 168	Nurses N = 90	Technologists N = 24
1. Sex, N (%)	Male	166, 59%	144, 86%	8, 9%	14, 58%
	Female	116, 41%	24, 14%	82, 91%	10, 42%
2. Age group, N (%)	20–29	45, 16%	17, 10%	18, 20%	10, 42%
	30–39	105, 37%	69, 41%	27, 30%	9, 38%
	40–49	103, 37%	71, 42%	29, 32%	3, 13%
	50–59	26, 9%	10, 6%	14, 16%	2, 8%
	60 and over	3, 1%	1, 1%	2, 2%	0, 0%
3. Job title		282	168, 60%	90, 32%	24, 9%
4. Institution size	University hospital or medical center	135, 48%	86, 51%	41, 46%	8, 33%
	Regional general hospital (>300 beds)	110, 39%	17, 10%	32, 36%	13, 54%
	Other	37, 13%	65, 39%	17, 19%	3, 13%
5. Career experience, years	1–5	45, 16%	24, 14%	12, 13%	9, 38%
	6–10	68, 24%	40, 24%	20, 22%	8, 33%
	11–15	56, 20%	37, 22%	16, 18%	3, 13%
	16–20	57, 20%	40, 24%	16, 18%	1, 4%
	Over 21	56, 20%	27, 16%	26, 29%	3, 13%
6. Operation of the fluoroscopy unit	Yes	188, 67%	156, 93%	23, 26%	9, 38%
7. Use of lead apron	Yes	281, 99%	168, 100%	89, 99%	24, 100%
8. Use of thyroid collar	Yes	90, 32%	46, 27%	35, 39%	9, 38%
9. Use of lead glasses	Yes	59, 21%	35, 21%	18, 20%	6, 25%
10. Use of radiation dosimeter	Yes	194, 69%	87, 52%	85, 94%	22, 92%
11. Fluoroscopy unit type	I don't know	30, 11%	11, 7%	17, 19%	2, 8%
12. RE of each procedure	Yes	42, 15%	21, 13%	12, 13%	9, 38%
13. Basic lecture on RE	Yes	180, 64%	119, 71%	49, 54%	12, 50%
14. Three principles of RP	Yes	167, 59%	102, 61%	53, 59%	12, 50%

Abbreviations: RE, radiation exposure; RP, radiation protection.

were aware of the three principles of radiation protection (Table 2)

Differences according to job title

Of the medical doctors, one hundred fifty-six (93%) had directly operated fluoroscopy units. By contrast, fewer nurses (N = 23, 26%) and technologists (N = 9, 38%) had operated fluoroscopy units because nurses and technologists are not licensed to manipulate radiation equipment except when instructed by an attending physician or dentist in Japan. The rates of wearing a lead apron among medical doctors, nurses, and technologists were 100% (N = 168), 99% (N = 89), and 100% (N = 24), respectively. Compared with the rates of wearing a lead apron, the rates of wearing a thyroid collar were low among medical doctors (27%, N = 46), nurses (39%, N = 35), and technologists (38%, N = 9).

Similarly, the rates of wearing lead glasses were also low among in medical doctors (21%, N = 35), nurses (20%, N = 18), and technologists (25%, N = 6). Rates of wearing radiation dosimeters were 52% (N = 87) by medical doctors, 94% (N = 85) by nurses, and 92% (N = 22) by technologists (Figure 1). Medical doctors were significantly less likely to wear dosimeters than the other medical workers ($p < 0.0001$). Regarding knowledge of radiation protection, 11 medical doctors (7%), 17 nurses (19%), and two technologists (8%) were not aware of the type of fluoroscopy unit. In total, 21 medical doctors (13%), 12 nurses (13%), and nine technologists (38%) were aware of the radiation dose of each procedure. One hundred nineteen medical doctors (71%), 49 nurses (54%), and 12 technologists (50%) had received lectures on radiation protection. In total, 102 medical doctors (61%), 53 nurses (59%), and 12 technologists (50%) were aware of the three principles of radiation protection.




FIGURE 1 Rates of use of protective equipment for radiation exposure according to job title

Factors affecting questionnaire answers

Use of equipment for radiation protection, including a lead apron, a thyroid collar, lead glasses, and a radiation dosimeter, did not differ based on years of experience. Similarly, the subjects did not differ in their recognition of the types of fluoroscopy; awareness of the radiation exposure dose of each procedure; attendance at basic lectures on radiation protection, including on the three principles of radiation protection; and size of the facility. However, medical doctors who were aware of the radiation exposure dose of each procedure wore dosimeters significantly more than those who were not ($p = 0.0008$). Medical doctors who had received basic lectures on radiation exposure were significantly more aware of the three principles of radiation protection ($p < 0.0001$), and those who were aware of the three principles were significantly more likely to have received lectures on radiation exposure ($p < 0.0001$).

DISCUSSION

This large Japanese questionnaire survey revealed that most of the medical staff in endoscopy departments usually wore lead aprons (almost 100%) but that they did not always wear thyroid collars (27%), lead glasses (21%), or dosimeters (52%) (Table 2, Figure 1). A previous Korean questionnaire survey conducted in 2011 showed similar adherence rates of endoscopists (aprons: 100%, thyroid collars: 52%, lead glasses: 14%, dosimeters 10%).⁸ Our cohort had slightly better rates of wearing dosimeters, but the rates were still insufficient. A recent document by the Ministry of Health, Labor, and Welfare of Japan reported that endoscopists had a lower dosimeter equipment usage rate than interventional radiologists (43% vs 100%).¹¹ Moreover, the rate of the usage of lead glasses by endoscopists in the current study was extremely low (21%). The authors of the

previous study reported that medical doctors and nurses in the gastroenterology department had much higher doses of radiation exposure to the lens than those of other departments, in which 11% of doctors and 50% of nurses received radiation exposure amounting to greater than 20 mSv/year.¹¹ Above all, the lack of knowledge of lens exposure causes a lower adherence rate for the use of lead glasses. There is an urgent necessity to gather information about global efforts, such as the revision of the upper limit of lens exposure doses in recent years.¹² The significant difference between the fields of radiology and cardiology and the field of gastroenterology may depend on the provision of academic lectures led by academic societies.

The rate of dosimeter use was 69% among all subjects. However, the rate of wearing a dosimeter among medical doctors (52%) was much lower than among nurses (92%) and technologists (94%), which led to lower overall adherence (Table 2, Figure 1). Campbell et al reported in 2002 that 47% of endoscopists performing ERCP never used dosimeters.¹³ Similarly, a Korean study from 2011 reported a rate of dosimeter use of 10%.⁸ This previous study showed that poor adherence has not improved for nearly two decades, which may reflect the situation in Japan, even though the previous survey was conducted in a different country. Radiation monitoring is also essential for basic fluoroscopic guidance practices that must be conducted to minimize exposure doses. However, only 21 doctors (13%) were aware of the radiation exposure dose of each procedure in the present survey.

The ICRP has stated the importance of radiation protection knowledge and education.⁴ However, there are still many reports of doctors' low awareness of radiation protection.¹⁴ Dauda et al reported that 80% of doctors had not attended basic lectures about radiation protection.¹⁵ Sethi et al conducted a questionnaire survey of 159 endoscopists in the United States and reported that the majority of endoscopists (62%) directly performed fluoroscopy during ERCP but that 57% had not received lectures on operating fluoroscopy equipment.⁹ Similarly, we believe that it is problematic that among the greater than 90% of the endoscopists in the present survey who operated fluoroscopy equipment, as only 71% took the basic courses. There have been many reports that education is useful for reducing radiation exposure. Specifically in the field of pediatric computed tomography, education and training programs for radiological institutes were concluded to be effective in achieving a substantial reduction in the radiation exposure dose.^{16–18} In the field of cardiovascular medicine, Georges et al demonstrated in 2009 that training in radiation protection for interventional cardiologists was associated with a 50% reduction in radiation exposure.¹⁹ In addition, sustained practice and X-ray system changes can result in a 40% decrease in radiation exposure.²⁰ In the field of gastroenterology,

TABLE 3 Factors affecting doctors' questionnaire responses

Questions	Years of experience	Size of facility	RE in each procedure	Basic lecture on RE	Three principles of RP
Lead apron (yes)	NSD	NSD	NSD	NSD	NSD
Thyroid collar (yes)	NSD	NSD	NSD	NSD	NSD
Lead glasses (yes)	NSD	NSD	NSD	NSD	NSD
Radiation dosimeter (yes)	NSD	NSD	$p = 0.0008$	NSD	NSD
Type of fluoroscopy unit (I don't know)	NSD	NSD	NSD	NSD	NSD
RE in each procedure (yes)	NSD	NSD	–	NSD	NSD
Basic lecture on RE (yes)	NSD	NSD	NSD	–	$p < 0.0001$
Three principles of RP (yes)	NSD	NSD	NSD	$p < 0.0001$	–

Abbreviations: RE, radiation exposure; RP, radiation protection; NSD, no significant difference.

Barakat et al reported the effectiveness of brief education for radiation protection.²¹ In the present survey, years of experience and the size of the facility did not affect the use of radiation protection, indicating that even experienced staff in high-volume center had not received enough education (Table 3). Our recent study revealed that awareness and education might reduce radiation exposure.²² We believe that the solution is to create an environment where education is widely available to both experienced and novice medical staff in endoscopy units; for example, mandatory educational lectures at conferences, such as those of radiological societies, may be considered.

In conclusion, this nationwide multicenter questionnaire survey of 282 medical staff showed the current status of protective equipment usage, awareness, and education in endoscopy departments in Japan. At present, the low rate of dosimeter wearing among gastroenterologists is a major problem, and there may be a lack of education in the gastrointestinal field in Japan. Continuing education can solve these problems, and endoscopists must be aware of the importance of radiation protection to protect both patients and staff. In addition, we are currently conducting a prospective nationwide study in Japan (REX-GI study) to collect actual radiation exposure data during digestive endoscopy. After this study, we plan to conduct a second questionnaire study to survey changes over the period between the two surveys.

CONFLICT OF INTEREST

The authors report no conflict of interest. The authors alone are responsible for the content and writing of this paper.

AUTHOR CONTRIBUTIONS

Shiro Hayashi planned this study and was the main author of the paper. Mamoru Takenaka and Makoto

Hosono advised the plan of this study. Tsutomu Nishida supervised and revised this paper. Mamoru Takenaka, Hirofumi Kogure, Takayuki Yakushijin, Hirotsugu Maruyama, Yasuki Hori, Toshiyuki Yoshio, Kenji Ikezawa, Tadayuki Takagi, Satoshi Asai, Kazuhiro Matsunaga, Kengo Matsumoto, Hidetaka Tsumura, Shinjiro Yamaguchi, Tetsuya Sumiyoshi, Koji Nagaike, Yuzuru Tamaru, Kazuo Hara, Toshio Fujisawa, Ichiro Oda, Ken Ohnita, Motohiko Kato, Hiroko Nebiki, Tatsuya Mikami, Akihiro Nishihara, Satoshi Egawa, and Ryuki Minami participated in this study. Participants were recruited at the individual institutions.

ETHICS DECLARATIONS

All participants were informed about the study. After giving their informed consent by checking the agree box in the web-based questionnaire, the participants were enrolled in the study. The study was performed following the guidelines of the declaration of Helsinki. Ethical approval was not sought for the present study because of the anonymous questionnaire survey.

FUNDING INFORMATION

This research received clinical research grants from the Japanese Society of Gastroenterology.

ORCID

Shiro Hayashi  <https://orcid.org/0000-0003-4533-2976>

Hirotsugu Maruyama  <https://orcid.org/0000-0002-8728-8101>

Toshiyuki Yoshio  <https://orcid.org/0000-0002-6546-0329>

Tetsuya Sumiyoshi  <https://orcid.org/0000-0002-9390-8477>

Tsutomu Nishida  <https://orcid.org/0000-0003-4037-9003>

REFERENCES

- De González AB, Darby S. Risk of cancer from diagnostic X-rays: estimates for the UK and 14 other countries. *The Lancet*. 2004;363: 345–51.
- Mathews JD, Forsythe AV, Brady Z, Butler MW, Goergen SK, Byrnes GB, et al. Cancer risk in 680,000 people exposed to computed tomography scans in childhood or adolescence: data linkage study of 11 million Australians. *BMJ*. 2013;346:f2360.
- ICRP. Avoidance of radiation injuries from medical interventional procedures ICRP publication 85. *Ann ICRP*. 2000;30:7–67.
- ICRP. Education and training in radiological protection for diagnostic and interventional procedures ICRP Publication 113. *Ann ICRP*. 2009;39:7–68.
- World Gastroenterology Organisation. Radiation protection in the endoscopy suite. 2020. <https://www.worldgastroenterology.org/guidelines/global-guidelines/radiation-protection-in-the-endoscopy-suite/radiation-protection-in-the-endoscopy-suite-english>. Accessed October 2, 2020.
- Dumonceau JM, Garcia-Fernandez FJ, Verdun FR, Carinou E, Donadille L, Damilakis J, et al. Radiation protection in digestive endoscopy: European Society of Digestive Endoscopy (ESGE) guideline. *Endoscopy*. 2012;44:408–21.
- Soye JA, Paterson A. A survey of awareness of radiation dose among health professionals in Northern Ireland. *Br J Radiol*. 2008;81:725–9.
- Son BK, Lee KT, Kim JS, Lee SO. Lack of radiation protection for endoscopists performing endoscopic retrograde cholangiopancreatography. *Korean J Gastroenterol*. 2011;58:93.
- Sethi S, Barakat MT, Friedland S, Banerjee S. Radiation training, radiation protection, and fluoroscopy utilization practices among US therapeutic endoscopists. *Dig Dis Sci*. 2019;64: 2455–66.
- Nishida T, Hayashi S, Takenaka M, Hosono M, Kogure H, Hasatani K, et al. Multicentre prospective observational study protocol for radiation exposure from gastrointestinal fluoroscopic procedures (REX-GI study). *BMJ Open*. 2020;10:e033604.
- Ministry of Health Labour and Welfare of Japan. [Japanese]. 2020. <https://www.mhlw.go.jp/content/11303000/000550109.pdf>. Accessed October 2, 2020.
- Ko S, Kang S, Ha M, Kim J, Jun JK, Kong KA, et al. Health effects from occupational radiation exposure among fluoroscopy-guided interventional medical workers: a systematic review. *J Vasc Interv Radiol*. 2018;29:353–66.
- Campbell N, Sparrow K, Fortier M, Ponich T. Practical radiation safety and protection for the endoscopist during ERCP. *Gastrointest Endosc*. 2002;55:552–7.
- Thomas KE, Parnell-Parmley JE, Haidar S, Moineddin R, Charkot E, Bendavid G, et al. Assessment of radiation dose awareness among pediatricians. *Pediatr Radiol*. 2006;36: 823–32.
- Dauda AM, Ozoh JO, Towobola OA. Medical doctors' awareness of radiation exposure in diagnostic radiology investigations in a South African academic institution. *SA J Radiol*. 2019;23:1707.
- Hojreh A, Weber M, Homolka P. Effect of staff training on radiation dose in pediatric CT. *Eur J Radiol*. 2015;84:1574–8.
- Schindera ST, Treier R, Von Allmen G, Nauer C, Trueb PR, Vock P, et al. An education and training programme for radiological institutes: impact on the reduction of the CT radiation dose. *Eur Radiol*. 2011;21:2039–45.
- Paolicchi F, Faggioni L, Bastiani L, Molinaro S, Caramella D, Bartolozzi C. Real practice radiation dose and dosimetric impact of radiological staff training in body CT examinations. *Insights Imaging*. 2013;4:239–44.
- Georges JL, Livarek B, Gibault-Genty G, Aziza JP, Hautecoeur JL, Soleille H, et al. Reduction of radiation delivered to patients undergoing invasive coronary procedures. Effect of a programme for dose reduction based on radiation-protection training. *Arch Cardiovasc Dis*. 2009;102:821–7.
- Fetterly KA, Mathew V, Lennon R, Bell MR, Holmes DR, Rihal CS. Radiation dose reduction in the invasive cardiovascular laboratory: implementing a culture and philosophy of radiation safety. *JACC Cardiovasc Interv*. 2012;5:866–73.
- Barakat MT, Thosani NC, Huang RJ, Choudhary A, Kochar R, Kothari S, et al. Effects of a brief educational program on optimization of fluoroscopy to minimize radiation exposure during endoscopic retrograde cholangiopancreatography. *Clin Gastroenterol Hepatol*. 2018;16:550–7.
- Hayashi S, Nishida T, Osugi N, Yamaoka S, Sugimoto A, Mukai K, et al. Time trend of the radiation exposure dose in endoscopic retrograde cholangiopancreatography over an 8-year period: a single-center retrospective study. *Am J Gastroenterol*. 2020;116:100–105.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Hayashi S, Takenaka M, Kogure H, et al. A questionnaire survey on radiation protection among 282 medical staff from 26 endoscopy-fluoroscopy departments in Japan. *DEN Open*. 2021;1:e5. <https://doi.org/10.1002/deo2.5>