



ARTICLE

Long-term maintenance of lung resident memory T cells is mediated by persistent antigen

Ida Uddbäck¹, Emily K. Cartwright², Amalie S. Schøller¹, Alexander N. Wein², Sarah L. Hayward², Jenna Lobby², Shiki Takamura³, Allan R. Thomsen¹, Jacob E. Kohlmeier^{2,4} and Jan P. Christensen¹

Tissue-resident memory T cells (T_{RM}) in the lungs are pivotal for protection against repeated infection with respiratory viruses. However, the gradual loss of these cells over time and the associated decline in clinical protection represent a serious limit in the development of efficient T cell based vaccines against respiratory pathogens. Here, using an adenovirus expressing influenza nucleoprotein (AdNP), we show that CD8 T_{RM} in the lungs can be maintained for at least 1 year post vaccination. Our results reveal that lung T_{RM} continued to proliferate in situ 8 months after AdNP vaccination. Importantly, this required airway vaccination and antigen persistence in the lung, as non-respiratory routes of vaccination failed to support long-term lung T_{RM} maintenance. In addition, parabiosis experiments show that in AdNP vaccinated mice, the lung T_{RM} pool is also sustained by continual replenishment from circulating memory CD8 T cells that differentiate into lung T_{RM} , a phenomenon not observed in influenza-infected parabiont partners. Concluding, these results demonstrate key requirements for long-lived cellular immunity to influenza virus, knowledge that could be utilized in future vaccine design.

Mucosal Immunology _#####_; <https://doi.org/10.1038/s41385-020-0309-3>

INTRODUCTION

CD8 tissue-resident memory T cells (T_{RM}) in the lungs represent a subset of memory T cells essential for optimal control of respiratory virus infections.^{1,2} Since T_{RM} were identified a decade ago, they have been found in various tissues including gut, skin, lung, reproductive tract, liver, and brain.^{3–7} T_{RM} are cells that reside in non-lymphoid tissues and function as a first line of defense against secondary infections. In addition to their unique anatomic location, T_{RM} have transcriptional profiles distinct from both central and effector memory T cells (T_{CM} and T_{EM} , respectively).^{8–10} In peripheral tissues, CD8 T_{RM} cells are primarily identified by their expression of the tissue retention markers CD69 and CD103, combined with the exclusion of cells that stain with an intravascular label.¹¹ However, not all T_{RM} are created equal. In the skin, T_{RM} remain in the tissue long after infection and antigen clearance.¹² Conversely, lung T_{RM} are lost a few months after an acute infection, leading to the loss of protection from secondary challenge.^{1,13} The reason for this rapid decay is still under investigation, but a harsh environment in the lungs and airways is likely to contribute. Recently published evidence shows that the airway environment causes transcriptional and epigenetic changes in the memory T cells resulting from amino acid starvation and leading to increased apoptosis.¹⁴ Previous reports have suggested residual antigen is important for continued development of virus-specific CD8 T cells after viral clearance following influenza infection.¹⁵ However, the potential role of residual antigen for the differentiation and maintenance of lung T_{RM} has not been investigated.

The discovery of the protective capacity of CD8 lung T_{RM} is of the utmost interest to the global vaccine community and, in particular, influenza vaccine research. However, the gradual waning of protection over time represents a serious limitation to the practical application of this finding. Therefore, a full understanding of the requirements for long-term lung T_{RM} maintenance would allow informed vaccine design to induce long-standing protective cellular immunity. Several studies have shown that adeno-based (Ad) vectors were effective at inducing CD8 T-cell-mediated protection against influenza.^{16–19} In addition to this, Ad vectors have been investigated as vaccines for both cancer and infections such as yellow fever, malaria, and *Listeria monocytogenes*.^{20–23} There are many advantages to the use of Ad vectors; they are easy to produce in high titers, have the potential to express large inserts and the vector itself functions as an efficient adjuvant.^{24,25} The possibility to enhance the immune response induced by Ad vectors further has been under investigation. One approach involves encoding of signals within the Ad vector to provide additional help to the CD8⁺ T-cell response.^{26,27} Previous work also investigated how the route of vaccination impacts the duration of protection against influenza virus. Mice vaccinated both intranasally (i.n.) and subcutaneously (s.c.) with an Ad vector expressing influenza nucleoprotein (AdNP) were protected from challenge longer than mice vaccinated by either the i.n. or s.c. route alone.¹⁸ However, these studies did not address the mechanism(s) of increased duration of protection, nor did they

¹Department of Immunology and Microbiology, University of Copenhagen, Copenhagen DK2200, Denmark; ²Department of Microbiology and Immunology, Emory University School of Medicine, Atlanta, GA 30322, USA; ³Department of Immunology, Faculty of Medicine, Kindai University, Osaka-Sayama, Osaka 589-8511, Japan and ⁴Emory-UGA Center of Excellence for Influenza Research and Surveillance, Atlanta, GA, USA

Correspondence: Jacob E. Kohlmeier (jkohlmeier@emory.edu) or Jan P. Christensen (jpc@sund.ku.dk)

These authors contributed equally: Ida Uddbäck, Emily K. Cartwright

These authors jointly supervised this work: Jacob E. Kohlmeier, Jan P. Christensen

Received: 13 September 2019 Revised: 20 April 2020 Accepted: 10 May 2020

Published online: 09 June 2020



address the impact of AdNP vaccination on the development and longevity of CD8⁺ T_{RM} in the lung and airways.

In the present study, we highlight the connection between persistent antigen expression following AdNP vaccination and maintenance of lung CD8 T_{RM}. We find, in contrast to the rapid loss of lung T_{RM} following influenza infection, mice primed with AdNP maintain a substantially larger population of lung T_{RM} for up to 1 year post vaccination (p.v.). Using Nur77^{GFP} reporter mice, we show lung T_{RM} of AdNP immunized mice continue to interact with cognate antigen for at least 3 months p.v. The expanded lung T_{RM} population in AdNP immunized mice is maintained in the lungs and airways by both in situ proliferation and continual replenishment of the lung T_{RM} pool from the circulation. Finally, we show that i.n. administration of AdNP is critical for the increased longevity of lung and airway T_{RM}. These results highlight the potential benefits of Ad vector vaccination, and underscore the importance of prolonged antigen expression in the lung for extended T-cell mediated protection against respiratory infections.

RESULTS

AdNP induced antigen-specific T_{RM} are maintained long-term in lung and BAL

To investigate the mechanism(s) underlying the increased duration of protection in AdNP immunized mice, we first examined the longevity of the CD8 T cells induced by AdNP and compared it to that generated by Influenza A/HKx31 (x31). After priming with either AdNP or x31, spleen, bronchoalveolar lavage (BAL), and lungs were isolated at various time points and numbers of D^bNP₃₆₆ tetramer-positive cells were determined by flow cytometry. For analysis of lung T_{RM} cells, intravital labeling was performed prior to exsanguination, labeling all cells within the circulation, ensuring that we could discriminate lung T_{RM} cells from cells in the lung vasculature.²⁸ Already at day 30 p.v. (Fig. 1a), there is a substantially larger population of D^bNP₃₆₆⁺ T cells in the lungs and airways of AdNP immunized mice. At day 180 (Fig. 1a), the proportion of D^bNP₃₆₆⁺ cells in lungs and airways of AdNP primed mice remain at the same level, or higher, whereas in x31 infected mice, the proportion of D^bNP₃₆₆⁺ cells declines substantially. Importantly, comparing the number of D^bNP₃₆₆⁺ cells between AdNP and x31 primed mice, we see that, despite that both generate high numbers in the acute phase (day 10–14), the numbers diverge as early as day 30 in lungs and airways (Fig. 1b). Where numbers of D^bNP₃₆₆⁺ cells in x31 infected mice rapidly decline, numbers are maintained in AdNP vaccinated mice up to at least day 275 p.v. In addition, a greater proportion of D^bNP₃₆₆⁺ cells express both CD69 and CD103 in the AdNP mice as early as day 30 p.v. (Fig. 1c). Even more striking, by day 90 (Fig. 1c) the proportion of CD69⁺CD103⁺ cells is maintained at around thirty percent in AdNP mice that declines to about five percent x31 infected mice. Examining AdNP mice at day 210, there is a further enrichment of CD69⁺CD103⁺ cells within the D^bNP₃₆₆⁺ population (Fig. 1c). A similar trend is observed when analyzing the absolute number (Fig. 1d), with the number of cells in the lungs and airways of AdNP mice remaining stable up to day 210 p.v. In contrast, numbers of CD69⁺CD103⁺ cells in the x31 primed mice are reduced drastically after the acute phase (Fig. 1d, e) and this population is almost completely lost by day 90. In addition to these data, a substantial number of D^bNP₃₆₆⁺ cells were still found in the lung and airways 580 days p.v., further illustrating the long-lived maintenance of the population in AdNP vaccinated mice (Fig. S1). The protective capacity of the AdNP induced CD8 T-cell response long after immunization have been previously demonstrated.¹⁸ Furthermore, through knock out and depletion experiments, we previously showed that the NP-specific CD8 T cells are responsible for the protective immunity established by AdNP vaccination. Importantly, we confirmed that the cellular immunity induced by AdNP vaccination was protective more than 255 days

p.v. (Fig. S2). Taken together, these data indicate that the long-lasting immunity induced by AdNP is due to increased duration of CD8 T_{RM} in the lung and airways.

Antigen persists in the lungs and airways after AdNP vaccination
Several studies have investigated the effect of persistent antigen on the CD8 T cell population following vaccination with adenoviral vectors.^{29,30} It is well established that low-level persistent antigen can stimulate T cells without causing exhaustion. This phenomenon, known as memory inflation has been studied in both murine *Cytomegalovirus* and Ad infections.^{29,31–33} Moreover, previous work has shown that antigen encounter in the lung is required for establishment of CD8 T_{RM}.^{34–36} Importantly, genomic material from adenoviruses and adeno-vectors persists in the tissue after immunization, in some cases up to a year post injection.^{30,37} In the present study, we confirmed persistence of the NP antigen by immunofluorescence microscopy in lungs 110 days after AdNP vaccination (Fig. S4). However, the effect of this low-level antigen persistence on lung T_{RM} has never been studied. To investigate this, we utilized a Nur77^{GFP} reporter mouse to visualize antigen-dependent CD8 T cell stimulation in vivo. The Nur77-GFP signal is transient and quickly lost after withdrawal of antigen stimulation, making the readout of GFP in Nur77^{GFP} mice a useful tool to investigate if lung T_{RM} have recently encountered their cognate antigen. Nur77^{GFP} mice were primed with x31 or AdNP and 45 or 90 days later BAL, lungs, and spleens were isolated and Nur77 expression analyzed. As illustrated, there is a significantly higher proportion of Nur77⁺CD69⁺ cells among the D^bNP₃₆₆⁺ CD8 T cells in the lungs and airways of AdNP mice at day 45 (Fig. 2a, b), compared with x31 mice. The frequency of Nur77⁺CD69⁺ decreases in both groups at day 90, however, we still find significantly more Nur77⁺CD69⁺ T cells in the lung T_{RM} of AdNP vaccinated mice. There is no difference in the frequency of Nur77⁺ cells between x31 and AdNP mice when expression was analyzed based on T_{RM} expression of CD69 and CD103 (data not shown).

Chronic PD-1 expression is believed to reflect ongoing antigen stimulation in T_{RM} populations^{1,38}; therefore, we examined PD-1 expression on CD8 lung T_{RM} as an additional indicator of persistent antigen stimulation (Fig. 2c, d). The majority of the D^bNP₃₆₆⁺ T cells in the BAL and lung of AdNP primed mice expressed PD-1 at day 60 and the proportion of CD69⁺PD-1⁺ cells was significantly higher in lungs and BAL of AdNP vaccinated mice compared with cells from x31 primed mice. PD-1 expression in AdNP mice was maintained until the last time point studied (day 275, data not shown). Since prolonged PD-1 expression can indicate a state of T-cell exhaustion,³⁹ it is important to stress that CD8 T_{RM} in AdNP vaccinated mice still show protection from PR8 challenge up to 255 days p.v. (Fig. S2). To directly address possible concerns regarding the functional relevance of PD-1⁺ expression on the airway T cells, the cellular response and its protective capacities after AdNP immunization were evaluated in PD-1^{-/-} mice (Fig. S3). We found no significant difference in the numbers of D^bNP₃₆₆⁺ T_{RM}, nor was there a significant difference in viral titers in the lungs 5 days after PR8 challenge compared with wild-type mice. We also examined TIM-3 expression, as TIM-3 has also been implicated in an exhausted T-cell phenotype.⁴⁰ D^bNP₃₆₆⁺ cells in AdNP immunized mice had little to no expression of TIM-3 (data not shown).

Next, we hypothesized that the persistent antigen in AdNP immunized mice results in continual in situ proliferation and thereby maintains the T_{RM} population. To evaluate proliferation, we administered EdU in the drinking water for a period of 7 days in both AdNP and x31 primed mice. Both at day 45 and day 90, mice immunized with AdNP have a significantly higher proportion of EdU⁺ cells in the BAL and lungs compared with x31 infected mice (Fig. 2e, f). Incredibly, we found EdU incorporation in D^bNP₃₆₆⁺ T cells of AdNP immunized mice as far out as 270 days p.v., with significantly greater incorporation in the BAL and lungs




Fig. 1 AdNP induced antigen-specific T_{RM} are maintained long-term in lung and BAL. C57BL/6 mice were immunized with AdNP subcutaneous (s.c.) in the footpad and i.n. or with HKx31 (x31) Influenza i.n. Lung, BAL, and Spleen were isolated for D^bNP_{366} tetramer analysis. **a** Representative plots. **b** Kinetics of absolute number of $D^bNP_{366}^+$ CD8 T cells. **c** Representative plots of residency markers CD69 and CD103 and **d** absolute numbers of $CD69^+CD103^+D^bNP_{366}^+$ CD8 T cells. **e** Fold reduction of $CD69^+CD103^+$ T cells. (**b + d**) Dots and bars represent mean and SD. **e** Dots represent fold reduction for each time point. All-time points are representative of three individual experiments with 5 mice in each.

(Fig. 2e, f). Due to the low number of $D^bNP_{366}^+$ T cells in x31 infected mice after day 90, we did not investigate EdU incorporation in these mice at later time points. Moreover, we observed that all lung T_{RM} subsets based on expression of CD69 and CD103 expression had a significantly higher proportion of EdU^+ NP-specific cells in AdNP immunized mice than observed in x31 infected mice (Fig. 2g, h). Taken together with the previous results, these data indicate the persistence of local antigen in the lung is driving activation and continued proliferation of lung T_{RM} following AdNP vaccination.

Persistent antigen in AdNP immunized mice pull circulating cells into the T_{RM} pool
It is well established that antigen is required for establishment and maintenance of lung T_{RM} following influenza infection.^{15,34,35}

However, we previously showed that i.n. vaccination with AdNP alone was not enough to induce long-lasting protection¹⁸ and hypothesized that the circulating population of NP-specific T cells induced by s.c. vaccination provides a pool of cells for continual recruitment and establishment of “new” T_{RM} following AdNP vaccination. To directly address this hypothesis, we utilized a parabiosis approach. Briefly, we immunized CD45 congenic mice with either AdNP or influenza x31 and 35 days later parabiotic surgery was performed. Twenty-eight days later, parabiotic partners were separated and spleen, lungs, and BAL were analyzed (Fig. 3a). Our data confirm a previous report showing that very few partner cells become resident in the BAL and lung of x31 infected mice.³⁵ Importantly, the proportion of donor cells found in the spleen is about 50%, indicating that equilibrium of recirculating memory cells was achieved (Fig. 3b, d). These data




Fig. 2 Antigen persists in the lungs and airways after AdNP vaccination. a C57BL/6xNur77^{GFP} mice were immunized with AdNP s.c.+i.n. or with x31 i.n. 45 or 90 days p.v., spleen, lung, and BAL were isolated for analysis of Nur77 expression. **a** Representative plots. **b** Percentages of Nur77⁺ cells within the D^bNP₃₆₆⁺ T cells. **c, d** C57BL/6 were immunized with AdNP i.n.+s.c. and 60 days later cell were isolated from BAL, lung and spleen and PD-1 and CD69 expression was analysed in the D^bNP₃₆₆ tetramer+ T cells. **c** Representative plots (**d**) percentage PD-1 and CD69 expression within the D^bNP₃₆₆ tetramer-positive population. **e-h** For proliferation studies, EdU incorporation was analysed in vaccinated C57BL/6 mice. 45, 90 and 270 days p.v., spleen, lung and BAL were isolated for analysis of EdU incorporation and CD44 in D^bNP₃₆₆⁺ T cells. **e** Representative plots. **f** Percentage of EdU⁺ out of D^bNP₃₆₆⁺ T cells (**g**) EdU incorporation in different subpopulations expressing CD69 and CD103. **h** Percentage of EdU in different CD69 and CD103 subpopulations. **p* < 0.05, ***p* < 0.01, The figures are representative of three individual experiments with 5 mice in each.




Fig. 3 Persistent antigen in AdNP immunized mice pull circulating cells into the T_{RM} pool. **a** AdNP and x31 primed mice were joined in a parabiosis surgery at early memory (5 weeks) and $D^{b}NP_{366}^{+}$ T cells were analysed after full equilibrium was reached 4 weeks after joining. **b** Representative plots of CD69, CD103 and CD49a expression in $D^{b}NP_{366}^{+}$ T cells in x31-primed partner (**c**) corresponding plots as in (**b**) for AdNP primed partner. **d + f** Percentage of host and donor cells in x31 primed partner (**e + g**) Percentage of host and donor in AdNP primed * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. The figure is representative of two individual experiments with 3 parabiosis pair in each.

are in stark contrast to parabiont partners immunized with AdNP, where AdNP had induced substantial recruitment of partner cells into all organ sites analyzed. Notably, in both BAL and lungs, about 50% of $D^{b}NP_{366}^{+}$ cells were from the x31-infected partner (Fig. 3c, e). In addition, partner cells in the lungs of AdNP vaccinated mice show similar expression of the T_{RM} markers CD69, CD103, and CD49a compared with host cells, indicating that the

recruited T cells differentiate into T_{RM} within the lungs. (Fig. 3c, right plots and 3g). In contrast, a very low proportion (about 1%) of the cells that had migrated from the partner into the lung and airways of the x31 infected mice expressed these tissue retention markers, indicating these are likely effector memory T cells transiting through the tissue (Fig. 3b right plots and 3f). These results provide compelling evidence that AdNP vaccination results in persistent antigen expression in the lungs of vaccinated mice, allowing circulating CD8 T cells to be continuously recruited into T_{RM} pool.

AdNP intranasal inoculation is indispensable for sustaining the lung T_{RM} pool

In order to further support our hypothesis of persistent local antigen expression as critical for maintenance of the lung T_{RM} population, we compared the NP-specific T-cell response in mice immunized both s.c. and i.n. with AdNP to mice vaccinated s.c. with AdNP and infected i.n. with x31 or mock-infected i.n. with PBS (Fig. 4a). By day 14 (Fig. 4b), mice immunized i.n. with AdNP had significantly more cells in the airway and lungs compared with mice inoculated i.n. with x31. At day 40 post priming, there were still significantly more $D^{b}NP_{366}^{+}$ T cells in in the airways and lungs of mice immunized with AdNP i.n. Importantly, regardless of s.c. administration of AdNP, the fold reduction of $D^{b}NP_{366}^{+}$ T cells in the lungs and airways between days 14 and 40 was greater in mice that received i.n. x31 or PBS (Fig. 4c). This emphasizes the necessity of local persistent antigen for the maintenance of a long-lived airway and lung T_{RM} population following AdNP vaccination.

DISCUSSION

Influenza virus infections represent a global health burden and currently available vaccines are inefficient. First, they are directed at a moving target, the main surface molecule, hemagglutinin, (HA), which is subject to substantial genetic variation as a result of both genetic drift as well as genetic shift. This creates a need for repeated vaccinations to sustain at least some protection. Second, the vaccine-induced immune response to HA consists largely of circulating IgG, whereas local immunity in the airways is limited. One way to remedy both of these deficiencies is to replace the current vaccine approach with one that induces a potent local T-cell response in the respiratory tract. Unlike antibodies, T cells also target the internal viral antigens, which are not subject to the same kind of selection as the surface molecules and therefore much more conserved between strains of influenza. However, a primary challenge in generating effective T cell based vaccines against respiratory pathogens, is the rapid decline of the lung T_{RM} population. However, our recently published results have indicated that combined local and systemic immunization in mice using AdNP induces a long-lasting protective CD8 population.¹⁸ Until now the underlying reasons for this sustained response has not been investigated. In this study, we provide evidence indicating that T_{RM} in AdNP vaccinated mice continue to encounter antigen in the lungs up to at least 3 months p.v., undergo in situ proliferation, and are continuously recruited into the lung from the circulating memory T cell pool.

First, our data provided evidence for long-standing proliferation of NP-specific cells in the lungs of Ad immunized mice in contrast to flu infected mice. Second, we observed ongoing recruitment of circulating memory cells as a consequence of prolonged antigen expression. Previous research has suggested that the T_{RM} population of influenza primed mice is maintained through a dynamic process consisting of a high apoptotic rate concurrent with replenishment from circulating memory cells, and that this recruitment is independent of local antigen.⁴¹ Based on our parabiosis experiments, we found little to no evidence of recruitment of circulating cells into the lung T_{RM} population in influenza primed mice. Moreover, a circulating memory T cell




Fig. 4 AdNP intranasal inoculation is indispensable for sustaining the lung T_{RM} pool. **a** C57BL/6 mice were immunized with AdNP s.c. +i.n. or, AdNP s.c.+x31 i.n. or, AdNP s.c.+PBS i.n. 14 and 40 days p.v. spleen, BAL and lungs were isolated and analysed for D^bNP₃₆₆⁺ tetramer cells. **b** Absolute numbers of D^bNP₃₆₆⁺ CD8 T cells in lung and BAL. **c** Fold reduction of D^bNP₃₆₆⁺ CD8 T cells in lung and BAL between day 14 and day 40. Dotted line represent 1 = no fold reduction. **p* < 0.05. The figure is representative of two individual experiments per time point with 5 mice in each.

population generated by s.c. AdNP immunization of i.n. influenza primed mice, did not suffice to maintain the lung T_{RM} population. Together, our data show the importance of prolonged local antigen expression for continuous recruitment of circulating T cells into the lung T_{RM} pool.

These data further emphasize that T_{RM} across tissues cannot be treated as one homogenous population, and that the requirements for establishment and maintenance are as diverse as the anatomic locations in which they reside. For example, antigen is not required for the establishment of skin T_{RM} and persistent antigen has not been shown to be necessary for their long-term maintenance.¹² Thus, it is highly plausible that long-term maintenance of a stable T_{RM} population in any given tissue is

dictated by how effectively local proliferation and ongoing recruitment from other memory cell subsets balances the local apoptotic rate.

Most importantly, all of these parameters are influenced by the microenvironment in the tissues of relevance. It should be acknowledged that other components of the immune system, such as APCs and even NP-specific antibodies, may also play a role in the maintenance of the CD8 T cell population.⁴² We have previously shown that CD8 T cells are the primary mechanism of protection after AdNP vaccination, but to what extent NP-specific antibodies are affected by the persisting antigen, and their contribution to protective immunity, remains to be investigated.¹⁸ In the lungs, the T_{RM} are subject to a relatively harsh environment, causing a high rate of apoptosis.¹⁴ While we cannot significantly change the environment of the lung, we have shown it is possible to expand and maintain the T_{RM} population by prolonged antigen stimulation. As stated previously, this strategy will open new doors for development of vaccines against flu and other viral respiratory diseases. In this context, it should be kept in mind that the dual vaccination approach applied for Ad immunized mice allows for antigen persistence in both the lungs and the periphery and both of these sources are likely contributing factors in the sustained response. It has been clearly documented that repeated antigen exposures gradually improve the quality of the primed cells, local cells become less prone to apoptosis, and circulating precursors develop a higher propensity to home to the lungs and differentiate into T_{RM}.^{43,44} The prolonged presence of antigen in the periphery is likely to have the same effect, in addition to increasing the number of activated circulating precursors. Consistent with this possibility, we see limited contraction following the initial T-cell burst in s.c. Ad immunized mice, and phenotypic analysis have revealed evidence of ongoing activation e.g. the prolonged-expression of CD43, on remaining antigen-specific CD8 T cells.^{45,46} The degree to which these factors, local antigen versus prolonged circulation of relevant precursors, contribute to the long-term immunity observed is not clear, but our previous results clearly show that combined s.c. and i.n. vaccination is superior to i. n. vaccination alone.¹⁸ Notably, due to the delicate nature of the lung, any vaccination strategy that results in persistent antigen must be carefully assessed for the induction of localized persistent inflammation and immunopathology that could be detrimental to the host. In conclusion, the results presented in this report show us that we can achieve a stable T-cell response in the lung with AdNP vaccination. Importantly, we also show that this long-lived T cell response is due to the presence of persistent local antigen. Not only is antigen required for T_{RM} formation as previously documented,³⁴ but unlike the situation in most other organ sites, persistent antigen is essential for the continual replenishment and long-term maintenance of T_{RM} population in the lungs.

METHODS

Experimental models

Six- to eight-week-old C57BL/6 mice from Taconic Biosciences were used in this study. All mice were rested upon arrival for at least 1 week. PepBoy/J, Nur77^{GFP} (Nur77-GFPCre B6-820), and PD-1^{-/-} (B6.Cg-Pdcd1tm1.1Shr/J) mice were purchased from Jackson Laboratory. All experimental procedures were approved by the national animal ethics committee (The Animal Experiments Inspectorate or IACUC) of the University of Copenhagen, Emory University, and Kindai University and were conducted in accordance with national guidelines; the mice were housed in an AAALAC accredited facility in accordance with good animal practice as defined by FELASA.

Virus and vaccines

All mice receiving intranasal (i.n.) inoculation were first anaesthetized by intraperitoneal (i.p.) injection with avertin (2,2,2

tribromoethanol in 2-methyl-2-butanol, 250 mg/kg). Influenza infection with A/Hong Kong/X31 (x31) was used at a dosage of 30,000 EID₅₀ in 30 µl HBSS and administered i.n. after avertin anesthesia. For influenza challenge, A/Puerto Rico/8/34 (PR8) was used at a dose of 3LD₅₀ in 30 µl i.n. The production of the replication-deficient adenovirus type 5 expressing influenza PR8 nucleoprotein (AdNP) used in this study has previously been described.¹⁸ Mice were immunized with 2 × 10⁷ plaque-forming units (PFU) in 30 µl of PBS i.n. and 30 µl s.c. in the right footpad after anesthesia with avertin.

Preparation of single-cell suspensions

To isolate resident lymphocytes in lung, mice were intravenously injected with 1.5 µg anti-CD3e [145-2C11] fluorophore-conjugated antibody in 200 µl PBS in the tail vein.^{11,28} Five minutes post injection, mice were anesthetized using avertin and exsanguinated. This was followed by harvest of BAL and other tissues.⁴⁷ After isolation lungs were digested with 5 g/L Collagenase D (Roche) and 2 × 10⁶ Units/L DNase (Sigma) for 30 min 37 °C. Samples were enriched by centrifugation in a 40%/80% Percoll gradient to isolate lymphocytes. Spleen and MLN were mechanically dissociated and passed through a 70 µm nylon filter prior to staining.

Antibodies for flow cytometry

Cells were first blocked for unspecific binding with αCD16/32 followed by staining with NP₃₆₆₋₃₇₄/Db tetramer conjugated to allophycocyanin (APC) or Brilliant Violet 421. Tetramer labeled cells was incubated with fluorophore-conjugated antibodies CD8α (clone 53-6.7), CD8β (clone H35 17.2) CD44 (clone IM7), CD45.1 (clone 30-F11), CD45.2 (clone 104) CD69 (clone H1.2F3), CD103 (clone 2E7), PD-1 and live/dead stain Zombie NIR. EdU staining was performed using Click It Plus Alexa Flour 647 Assay kit (Invitrogen) according to manufacturer's instructions. Samples were analyzed on a Fortessa LSR II (BD Biosciences). Data analysis was conducted using FlowJo v10 software (TreeStar). Gates for CD69, CD103, and PD-1 were set using fluorescence minus one samples. All antibodies were purchased from Biolegend. Relevant tetramers were kindly provided by Søren Buus, Department of Immunology and Microbiology and the NIH tetramer core facility.

Parabiosis

For parabiosis surgery, mice were anesthetized, and a clipper was used to remove flank hair. This was followed by a longitudinal skin incision from the knee to the elbow on a single lateral side along with a 1-cm lateral peritoneal incision. Suturing each reciprocal peritoneal opening joined the two mice together. To hold the mice in the upright position, two mattress stiches were made on the lateral edges of the skin section. Also, the dorsal and ventral sides of the skin section of each mouse were further joined with wound clips. Equilibrium was confirmed 10 days after surgery with a blood sample.

MDCK plaque assay

Lungs were homogenized in 1%BSA in PBS 9X the lung weight to obtain a 10% w/v suspension. The mixture was homogenized using sterilized sand, mortar, and pestle. This was followed by centrifugation at 600 × g, 15 min, 4 °C. The supernatant was transferred to a new tube and kept on ice until use. 4.5 × 10⁴ MDCK cells were seeded in 100 µl medium in 96-well plates and the following day, lung supernatant was added in tenfold dilutions in media containing DMEM 1965 medium with 2 mM L-glutamin, 200 IU/ml penicillin, 50 µg/ml streptomycin, 0.2% BSA, 1% sodium pyruvate and 5 units/ml TPCK Trypsin for 2 h. Virus was then removed and samples were incubated for 48 h, 37 °C, 5% CO₂, with a 1:1 mixture of medium containing 2× minimum essential medium (MEM) eagle supplemented with 0.4% BSA, 10% NaHCO₃, 2% Streptomycin, 2% penicillin and 5 units/ml TPCK trypsin and 1.8% methyl cellulose. After incubation overlay was removed and

cells were fixed with 4% formaldehyde in PBS for 30 min at room temperature (RT) and permeabilized using warm 0.5% Triton-X in Hanks balanced salt solution for 10 min at RT. After permeabilization cells were incubated with primary α-influenza nucleocapsid A mAb (Nordic Biosite) diluted 1:1500 in 10% FBS in PBS at 37 °C, 5% CO₂. Following the primary antibody, cells were incubated with secondary goat α-mouse HRP conjugated mAb (Dako) diluted 1:500 in 10% FBS in PBS at 37 °C. After the secondary antibody, substrate was added containing 3 mg/ml 3-amino-9-ethylcarbazole and 0.07% H₂O₂ and 5 mM citrate phosphate buffer pH5 and incubated at RT for 30 min. After these PFU were counted and calculated per g lung according to the following formula:

$$\text{Average \# of plaque/well} \times \text{Dilution factor} \times 20 = \text{PFU/g lung}$$

Immunofluorescence microscopy

For OCT imaging, mice were infected with the specified agent and harvested at indicated days. The mice were euthanized by 2,2,2-tribromoethanol overdose and the ribcage was dissected off. An incision was made in the trachea and the airways were inflated with 0.75 mL OCT via an 18-gauge iv catheter. The trachea was tied using 5-0 silk suture on a reverse cutting needle and the heart, lungs, and thymus were removed en bloc and flash frozen in OCT. Sections were cut at 7 µm and transferred to slides for staining. Slides were fixed in 75% acetone/25% ethanol solution for 10 min and blocked with 10% donkey serum, 10% mouse serum, 10% rat serum, 1 µg/mL anti CD16/32 clone 2.4G2, and 5% FCS in PBS for 30 min on ice. Slides were stained with primary antibodies for 30 min on ice and secondary reagents for 15 min on ice. Coverslips were placed with ProFade Gold mounting media and imaged using a Zeiss AxioObserver microscope using Zen 2 software. Antibodies used include anti-EpCAM-A647 (clone G8.8, Biologend), anti-Collagen-IV goat pAb (part # AB769, EMD Millipore), Donkey anti-goat-A405 (part # 705-475-147, Jackson ImmunoResearch), anti-influenza-nucleoprotein-FITC (clone 431, abcam), anti-FITC-A488 (part # A-11090, Invitrogen), and anti-CD11c-A594 (clone N418, Biologend).

Statistical analysis

Statistical analysis was performed using GraphPad Prism 7. ANOVA was used for statistical testing for all experiments and where pairwise comparison was made Mann-Whitney rank test was used. **p* < 0.05, ***p* < 0.01, ****p* < 0.001.

ACKNOWLEDGEMENTS

This project was supported by the Danish Research Council and 'Fonden til Lægevidenskabens Fremme' for grant support. IEMU is the recipient of a PhD scholarship from the Faculty of Health and Medical Sciences, University of Copenhagen. We would also like to acknowledge the support by NIH grants HL122559, HL138508, and Centers of Excellence in Influenza Research and Surveillance contract HHSN272201400004C (to J.E.K.), and Grant-in-Aid for Young Scientists (A) 24689043, Grant-in-Aid for Scientific Research (C) 16K08850 from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and grants from Takeda Science Foundation, Daiichi-Sankyo Foundation of Life Science, Uehara Memorial Foundation, and Kanae Foundation for Promotion of Medical Science (to S.T.). S.L.H. was supported by NIH grant F31 HL136101. We recognize contributions from the Children's Healthcare of Atlanta and Emory University Pediatric Flow Cytometry Core for cell sorting and the NIH Tetramer Core Facility (contract HHSN272201300006C).

AUTHOR CONTRIBUTIONS

I.U. and E.K.C. designed, performed, and analyzed most of the experiments with input from A.R.T., J.E.K., and J.P.C. A.S.S., S.L.H., J.L. carried out tetramer stainings and facs analysis. S.T. performed parabiosis experiments. A.N.W. designed and performed



immunofluorescence microscopy experiment and analysis. I.U. and E.K.C. wrote the manuscript and A.R.T., J.E.K., and J.P.C. edited the manuscript.

ADDITIONAL INFORMATION

The online version of this article (<https://doi.org/10.1038/s41385-020-0309-3>) contains supplementary material, which is available to authorized users.


Competing interests: The authors declare no competing interests.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.


REFERENCES

1. Wu, T. et al. Lung-resident memory CD8 T cells (TRM) are indispensable for optimal cross-protection against pulmonary virus infection. *J. Leukoc. Biol.* **95**, 215–224 (2014).
2. Kinnear, E. et al. Airway T cells protect against RSV infection in the absence of antibody. *Mucosal. Immunol.* **11**, 290 (2018).
3. Gebhardt, T. et al. Memory T cells in nonlymphoid tissue that provide enhanced local immunity during infection with herpes simplex virus. *Nat. Immunol.* **10**, 524–530 (2009).
4. Jiang, X. et al. Skin infection generates non-migratory memory CD8+ T(RM) cells providing global skin immunity. *Nature* **483**, 227–231 (2012).
5. Masopust, D. et al. Dynamic T cell migration program provides resident memory within intestinal epithelium. *J. Exp. Med.* **207**, 553–564 (2010).
6. Schenkel, J. M., Fraser, K. A., Vezys, V. & Masopust, D. Sensing and alarm function of resident memory CD8(+) T cells. *Nat. Immunol.* **14**, 509–513 (2013).
7. Stelma, F. et al. Human intrahepatic CD69 + CD8+ T cells have a tissue resident memory T cell phenotype with reduced cytolytic capacity. *Sci. Rep.* **7**, 6172 (2017).
8. Mackay, L. K. et al. T-box transcription factors combine with the cytokines TGF-beta and IL-15 to control tissue-resident memory T cell fate. *Immunity* **43**, 1101–1111 (2015).
9. Wakim, L. M. et al. The molecular signature of tissue resident memory CD8 T cells isolated from the brain. *J. Immunol.* **189**, 3462–3471 (2012).
10. Kumar, B. V. et al. Human tissue-resident memory T cells are defined by core transcriptional and functional signatures in lymphoid and mucosal sites. *Cell Rep.* **20**, 2921–2934 (2017).
11. Anderson, K. G. et al. Intravascular staining for discrimination of vascular and tissue leukocytes. *Nat. Protoc.* **9**, 209–222 (2014).
12. Mackay, L. K. et al. Long-lived epithelial immunity by tissue-resident memory T (TRM) cells in the absence of persisting local antigen presentation. *Proc. Natl Acad. Sci. USA* **109**, 7037–7042 (2012).
13. Gilchuk, P. et al. A distinct lung-interstitium-resident memory CD8(+) T cell subset confers enhanced protection to lower respiratory tract infection. *Cell Rep.* **16**, 1800–1809 (2016).
14. Hayward, S. L. et al. Environmental cues regulate epigenetic reprogramming of airway-resident memory CD8(+) T cells. *Nat. Immunol.* **21**, 309–320 (2020).
15. Zammit, D. J., Turner, D. L., Klonowski, K. D., Lefrancois, L. & Cauley, L. S. Residual antigen presentation after influenza virus infection affects CD8 T cell activation and migration. *Immunity* **24**, 439–449 (2006).
16. Coughlan, L. et al. Heterologous two-dose vaccination with simian adenovirus and poxvirus vectors elicits long-lasting cellular immunity to influenza virus in healthy adults. *EBioMed.* **29**, 146–154 (2018).
17. Powell, T. J. et al. Examination of influenza specific T cell responses after influenza virus challenge in individuals vaccinated with MVA-NP+M1 vaccine. *PLoS ONE* **8**, e62778 (2013).
18. Uddback, I. E. et al. Combined local and systemic immunization is essential for durable T-cell mediated heterosubtypic immunity against influenza A virus. *Sci. Rep.* **6**, 20137 (2016).
19. Vitelli, A. et al. Vaccination to conserved influenza antigens in mice using a novel Simian adenovirus vector, PanAd3, derived from the bonobo Pan paniscus. *PLoS ONE* **8**, e55435 (2013).
20. Bassi, M. R. et al. Vaccination with replication deficient adenovectors encoding YF-17D antigens induces long-lasting protection from severe yellow fever virus infection in mice. *PLoS Negl. Trop. Dis.* **10**, e0004464 (2016).
21. Jensen, S. et al. Adenovirus-based vaccine against *Listeria monocytogenes*: extending the concept of invariant chain linkage. *J. Immunol.* **191**, 4152–4164 (2013).


22. Lasaro, M. O. & Ertl, H. C. New insights on adenovirus as vaccine vectors. *Mol. Ther.* **17**, 1333–1339 (2009).
23. Sheehy, S. H. et al. Phase Ia clinical evaluation of the safety and immunogenicity of the Plasmodium falciparum blood-stage antigen AMA1 in ChAd63 and MVA vaccine vectors. *PLoS ONE* **7**, e31208 (2012).
24. Bett, A. J., Prevec, L. & Graham, F. L. Packaging capacity and stability of human adenovirus type 5 vectors. *J. Virol.* **67**, 5911–5921 (1993).
25. Kamen, A. & Henry, O. Development and optimization of an adenovirus production process. *J. Gene Med.* **6**(Suppl 1), S184–S192 (2004).
26. Zhou, A. C., Wagar, L. E., Wortzman, M. E. & Watts, T. H. Intrinsic 4-1BB signals are indispensable for the establishment of an influenza-specific tissue-resident memory CD8 T-cell population in the lung. *Mucosal. Immunol.* **10**, 1294–1309 (2017).
27. Jensen, B. A. et al. Co-expression of tumor antigen and interleukin-2 from an adenoviral vector augments the efficiency of therapeutic tumor vaccination. *Mol. Ther.* **22**, 2107–2117 (2014).
28. Anderson, K. G. et al. Cutting edge: intravascular staining redefines lung CD8 T cell responses. *J. Immunol.* **189**, 2702–2706 (2012).
29. Bolinger, B. et al. A new model for CD8+ T cell memory inflation based upon a recombinant adenoviral vector. *J. Immunol.* **190**, 4162–4174 (2013).
30. Tatsis, N. et al. Adenoviral vectors persist in vivo and maintain activated CD8+ T cells: implications for their use as vaccines. *Blood* **110**, 1916–1923 (2007).
31. Karrer, U. et al. Memory inflation: continuous accumulation of antiviral CD8+ T cells over time. *J. Immunol.* **170**, 2022–2029 (2003).
32. Holtappels, R., Pahl-Seibert, M. F., Thomas, D. & Reddehase, M. J. Enrichment of immediate-early 1 (m123/pp89) peptide-specific CD8 T cells in a pulmonary CD62L(lo) memory-effector cell pool during latent murine cytomegalovirus infection of the lungs. *J. Virol.* **74**, 11495–11503 (2000).
33. Holst, P. J., Orskov, C., Thomsen, A. R. & Christensen, J. P. Quality of the transgene-specific CD8+ T cell response induced by adenoviral vector immunization is critically influenced by virus dose and route of vaccination. *J. Immunol.* **184**, 4431–4439 (2010).
34. McMaster, S. R. et al. Pulmonary antigen encounter regulates the establishment of tissue-resident CD8 memory T cells in the lung airways and parenchyma. *Mucosal. Immunol.* **11**, 1071–1078 (2018).
35. Takamura, S. et al. Specific niches for lung-resident memory CD8+ T cells at the site of tissue regeneration enable CD69-independent maintenance. *J. Exp. Med.* **213**, 3057–3073 (2016).
36. Pizzolla, A. et al. Resident memory CD8+ T cells in the upper respiratory tract prevent pulmonary influenza virus infection. *Sci. Immunol.* **2**, eaam6970 (2017).
37. Finn, J. D. et al. Persistence of transgene expression influences CD8+ T-cell expansion and maintenance following immunization with recombinant adenovirus. *J. Virol.* **83**, 12027–12036 (2009).
38. Sakai, S. et al. Cutting edge: control of Mycobacterium tuberculosis infection by a subset of lung parenchyma-homing CD4 T cells. *J. Immunol.* **192**, 2965–2969 (2014).
39. Day, C. L. et al. PD-1 expression on HIV-specific T cells is associated with T-cell exhaustion and disease progression. *Nature* **443**, 350–354 (2006).
40. Blackburn, S. D. et al. Coregulation of CD8+ T cell exhaustion by multiple inhibitory receptors during chronic viral infection. *Nat. Immunol.* **10**, 29–37 (2009).
41. Slutter, B., et al. Dynamics of influenza-induced lung-resident memory T cells underlie waning heterosubtypic immunity. *Sci. Immunol.* **2** (2017).
42. Leon, B., Ballesteros-Tato, A., Randall, T. D. & Lund, F. E. Prolonged antigen presentation by immune complex-binding dendritic cells programs the proliferative capacity of memory CD8 T cells. *J. Exp. Med.* **211**, 1637–1655 (2014).
43. Van Braeckel-Budimir, N., Varga, S. M., Badovinac, V. P. & Harty, J. T. Repeated antigen exposure extends the durability of influenza-specific lung-resident memory CD8(+) T cells and heterosubtypic immunity. *Cell Rep.* **24**, 3374–3382 e3373 (2018).
44. Tatsis, N. et al. Multiple immunizations with adenovirus and MVA vectors improve CD8+ T cell functionality and mucosal homing. *Virology* **367**, 156–167 (2007).
45. Steffensen, M. A. et al. Qualitative and quantitative analysis of adenovirus type 5 vector-induced memory CD8 T cells: not as bad as their reputation. *J. Virol.* **87**, 6283–6295 (2013).
46. Jahn, M. L., Steffensen, M. A., Christensen, J. P. & Thomsen, A. R. Analysis of adenovirus-induced immunity to infection with *Listeria monocytogenes*: Fading protection coincides with declining CD8 T cell numbers and phenotypic changes. *Vaccine* **36**, 2825–2832 (2018).
47. McMaster, S. R., Wilson, J. J., Wang, H. & Kohlmeier, J. E. Airway-resident memory CD8 T cells provide antigen-specific protection against respiratory virus challenge through rapid IFN-gamma production. *J. Immunol.* **195**, 203–209 (2015).




Supplementary Figure 1. AdNP generate long lasting lung and airway populations. C57BL/6 mice were immunized with AdNP s.c. and i.n. and 580 days later organs were isolated and D^bNP₃₆₆ tetramer positive cells were enumerated through flow cytometry.



Supplementary Figure 2. AdNP generate long lasting protective immunity. C57BL/6 mice were immunized with AdNP s.c. and i.n. or with x31 Influenza i.n. 255 days later mice were challenged with PR8 and weightloss (**A**) was recorded and lung titer was determined 7 days after infection (**B**). Dots in (**A**) represents mean±SD. *:p<0.05, (**B**) line and bars represent mean±SD. Each dot represents one mouse. Dotted line represents detection limit of assay. PFU: plaque forming units



Supplementary Figure 3. AdNP vaccination in PD-1 knock out mice results in generation of similar CD8 T cell response. WT C57BL/6 mice and PD-1 knock out mice (PD-1^{-/-}) were immunized with AdNP s.c. and i.n. 60 days later BAL, lung, MLN and spleen was isolated and (A) D^bNP₃₆₆ tetramer positive cells were enumerated through flow cytometry. (B) Number of CD69 and CD103 subsets out of the tetramer positive T cells were also analysed. (C) 60 days after vaccination, groups of mice were challenged with PR8 and titers in lungs were determined five days after challenge. (A) Bars represents mean±SD, each dot represents one mouse. (B) Line and bars represent mean±SD. Each dot represents one mouse. Dotted line represents detection limit of assay, ns: no significant difference



Supplementary Figure 4: Influenza NP protein persists in the lungs long after AdNP vaccination.

Immunofluorescence microscopy of OCT-embedded mouse lungs for the influenza nucleoprotein following indicated infections. Colors are blue- Collagen IV, white- EpCAM, red- CD11c, green- FluNP. Scale bars indicate 50 um. Data are representative of two mice after AdNP vaccination.