

CANCELLATION OF FLUCTUATION IN STOCHASTIC RANKING PROCESS WITH SPACE-TIME DEPENDENT INTENSITIES

TETSUYA HATTORI*

Abstract

We consider the stochastic ranking process with space-time dependent unbounded jump rates for the particles. We prove that the joint empirical distribution of jump rate and scaled position converges almost surely to a deterministic distribution in the infinite particle limit. We assume topology of weak convergence for the space of distributions, which implies that the fluctuations among particles with different jump rates cancel in the limit. The results are proved by first finding an auxiliary stochastic ranking process, for which a strong law of large numbers is applied, and then applying a multi time recursive Gronwall's inequality. The limit has a representation in terms of non-Markovian processes which we call point processes with last-arrival-time dependent intensities. We also prove the propagation of chaos, i.e., the tagged particle processes also converge almost surely.

1 Introduction.

Let N be a positive integer and $T > 0$. T is an arbitrary constant fixed throughout the paper, and we are interested in the limit $N \rightarrow \infty$. A stochastic ranking process is a stochastic system of N particles on a line segment $[0, 1]$ for a time interval $[0, T]$, defined as follows. Let W be a set of non-negative valued continuous functions in two variables,

2010 *Mathematics Subject Classification*. Primary 60K35; Secondary 82C22.

Key words and phrases. Stochastic ranking process, hydrodynamic limit, law of large numbers, complete convergence, last-arrival-time dependent intensity, Gronwall inequality.

*This work is supported by JSPS KAKENHI Grant Number 26400146 from Japan Society for the Promotion of Science, and by Keio Gijuku Academic Development Funds from Keio University.

URL: <http://web.econ.keio.ac.jp/staff/hattori/research.htm>

continuously differentiable in the first variable, and let w_1, w_2, \dots be an infinite sequence in W .

Let $y_1^{(N)}, y_2^{(N)}, \dots, y_N^{(N)}$ be a permutation of $\{\frac{i}{N} \mid i = 0, 1, \dots, N-1\}$. Then a stochastic ranking process is a system of stochastic processes $\{Y_i^{(N)} \mid i = 1, 2, \dots, N\}$ defined on a probability space (Ω, \mathcal{F}, P) by

$$(1) \quad \begin{aligned} Y_i^{(N)}(t) &= y_i^{(N)} \\ &+ \frac{1}{N} \sum_{j=1}^N \int_{s \in (0, t]} \int_{\xi \in [0, \infty)} \mathbf{1}_{Y_j^{(N)}(s-) > Y_i^{(N)}(s-)} \mathbf{1}_{\xi \in [0, w_j(Y_j^{(N)}(s-), s))} \nu_j^{(N)}(d\xi ds) \\ &- \int_{s \in (0, t]} \int_{\xi \in [0, \infty)} Y_i^{(N)}(s-) \mathbf{1}_{\xi \in [0, w_i(Y_i^{(N)}(s-), s))} \nu_i^{(N)}(d\xi ds), \\ &i = 1, 2, \dots, N, \quad t \geq 0. \end{aligned}$$

Here, $\mathbf{1}_A$ is the indicator function of an event A , and for each N , $\nu_i^{(N)}$, $i = 1, 2, \dots, N$, are independent Poisson random measures on $[0, \infty) \times [0, \infty)$ with uniform unit intensity measures, i.e., $E[\nu_i^{(N)}([a, b] \times [c, d])] = (b-a)(d-c)$ for $b > a \geq 0$ and $d > c \geq 0$, and $\nu_i^{(N)}(A)$ and $\nu_i^{(N)}(B)$ are independent Poisson variables if $A \cap B = \emptyset$.

If we put

$$(2) \quad \tilde{\nu}_i^{(N)}(t) = \int_{s \in (0, t]} \int_{\xi \in [0, \infty)} \mathbf{1}_{\xi \in [0, w_i(Y_i^{(N)}(s-), s))} \nu_i^{(N)}(d\xi ds),$$

then (1), the position of the particle i at time t , is expressed as

$$Y_i^{(N)}(t) = y_i^{(N)} + \frac{1}{N} \sum_{j=1}^N \int_0^t \mathbf{1}_{Y_j^{(N)}(s-) > Y_i^{(N)}(s-)} \tilde{\nu}_j^{(N)}(ds) - \int_0^t Y_i^{(N)}(s-) \tilde{\nu}_i^{(N)}(ds).$$

We then see that the time development of $\{Y_i^{(N)}\}$ is determined by the move-to-front rules [18] driven by the point processes $\{\tilde{\nu}_i^{(N)}\}$ with space-time dependent intensities given by the ‘density functions’ $\{w_i\}$. If w_i is constant then the process $\tilde{\nu}_i^{(N)}$ is the Poisson process, while in general, its increments depend on past and is dependent on other particles. The move-to-front rule in particular implies that the particle system $\{Y_i^{(N)}\}$ as a whole takes values in the rearrangement of $\frac{i}{N}$, $i = 0, 1, \dots, N-1$. Each particle either increases its position by $\frac{1}{N}$, or else takes the value 0, i.e. jumps to the top position, as t increases.

A starting point for our study is the joint empirical distribution of w_i and the position, given by

$$(3) \quad \mu_t^{(N)} = \frac{1}{N} \sum_{i=1}^N \delta_{(w_i, Y_i^{(N)}(t))}.$$

Here δ_c is a unit measure concentrated at c . (We will use this notation for a unit measure on any probability space.) $\mu_t^{(N)}$, $t \in [0, T]$, is regarded as a stochastic process taking values in the set of Borel probability measures on $C^{1,0}([0, 1] \times [0, T]) \times [0, 1]$. Here $C^{1,0}([0, 1] \times [0, T])$ is the total set of functions $f \in C([0, 1] \times [0, T])$ such that $\frac{\partial f}{\partial y} \in C([0, 1] \times [0, T])$. Since $C^{1,0}([0, 1] \times [0, T])$ is a Polish space with norm

$$\sup_{(y,t) \in [0,1] \times [0,T]} \max \left\{ |w(y, t)|, \left| \frac{\partial w}{\partial y}(y, t) \right| \right\},$$

so is $C^{1,0}([0, 1] \times [0, T]) \times [0, 1]$ (see [1, Example 26.2]). We assume a standard topology of weak convergence of probability measures on $C^{0,1}([0, 1] \times [0, T]) \times [0, 1]$. In particular, the initial distribution is

$$(4) \quad \mu_0^{(N)} = \frac{1}{N} \sum_{i=1}^N \delta_{(w_i, y_i^{(N)})}.$$

By considering a process $X_i^{(N)}(t) = NY_i^{(N)}(t) + 1$ taking values in positive integers, we see that a stochastic ranking process is a model of ranking system, such as the sales ranks found at online bookstores[6, 7, 5, 8, 10, 9, 16, 17, 14]. As a model of popularity ranks of an online bookstore, the move-to-front rule defines the rank as a stochastic number with the ‘latest purchased book as most popular’ rule. That the intensity w_i differs for different book i represents that there are popular books and less popular ones. (Many books on mathematics perhaps provide examples of the latter.) See also [14] and references therein for more background.

We will prove existence of hydrodynamic limit, that assuming convergence of $\mu_0^{(N)}$ as $N \rightarrow \infty$ we have convergence of $\mu_t^{(N)}$ for all $t \in [0, T]$. The common standard quantities effective for the move-to-front rules are the characteristic curves $Y_C^{(N)}$ defined by

$$(5) \quad \begin{aligned} Y_C^{(N)}(\gamma, t) &= y_0 \\ &+ \frac{1}{N} \sum_{j=1}^N \int_{s \in (t_0, t]} \int_{\xi \in [0, \infty)} \mathbf{1}_{Y_j^{(N)}(s-) \geq Y_C^{(N)}(\gamma, s-)} \\ &\quad \times \mathbf{1}_{\xi \in [0, w_j(Y_j^{(N)}(s-), s))} \nu_j^{(N)}(d\xi ds), \\ \gamma &= (y_0, t_0) \in [0, 1] \times [0, T], \quad t \geq t_0, \end{aligned}$$

and a set of spatial distribution functions $\varphi^{(N)}$ of $\mu_t^{(N)}$ defined by

$$(6) \quad \begin{aligned} \varphi^{(N)}(dw, \gamma, t) &= \mu_t^{(N)}(dw \times [Y_C^{(N)}(\gamma, t), 1]), \\ \gamma &= (y_0, t_0) \in [0, 1] \times [0, T], \quad t \geq t_0, \quad w \in W. \end{aligned}$$

The latter is a refinement of the former in the sense that (3) and (6) imply

$$(7) \quad \begin{aligned} Y_C^{(N)}(\gamma, t) &= y_0 + \frac{[N(1 - y_0)]}{N} - \varphi^{(N)}(W, \gamma, t), \\ \gamma &= (y_0, t_0) \in [0, 1] \times [0, T], \quad t \geq t_0. \end{aligned}$$

Since $\varphi^{(N)}$ determines $\mu_t^{(N)}$, the convergence problem of $\mu_t^{(N)}$ is reduced to that of $\varphi^{(N)}$.

If the intensity densities $w \in W$ are *independent* of y , then $\varphi^{(N)}$ is an arithmetic mean of *independent* stochastic processes, so that a strong law of large numbers for a sum of *independent processes* can be applied to prove existence of almost sure $N \rightarrow \infty$ limit under reasonable assumptions [5]. In contrast, when the intensity densities depend on y , the problem is a more involved one of law of large numbers for *dependent processes*. The case of spatially varying intensity densities was first studied in [14], where we found the existence of the limit under restricting assumptions. Firstly, we assumed in [14] that $w \in W$ is bounded as well as the derivative in the first variable. We keep the latter assumption

$$(8) \quad C_W := \sup_{w \in W} \sup_{(y,t) \in [0,1] \times [0,T]} \left| \frac{\partial w}{\partial y}(y, t) \right| < \infty$$

in the present study, but replace the assumption of boundedness of w in [14] by a milder (and natural, in view of the previous study [5] for simpler cases) assumption (24). This improvement in the choice of assumption in particular allows Pareto (power law) distributions for λ in (24) (which was excluded in [14]), which may be of interest in applying stochastic ranking processes to social studies [9, 10]. Secondly, we adopted the total variation norm for the topology of the space of Borel measures in [14]. This means that, among other points, the law of large numbers proved in [14] is a cancellation of fluctuations among processes $\{Y_i^{(N)}\}$ having the same associated intensity densities $w_i = w$. This assumption is to be compared with the results in [5], where we proved a corresponding convergence theorem (for the easier case of spatially constant intensity densities) with topology of the space of Borel measures induced by weak convergence, which, for example, allows all the w_i 's to be different and the cancellation of fluctuation is still implied. It would be mathematically interesting to see how this law of large numbers (fluctuation cancellation) mechanism, which escaped from our hands in [14], would be stable against introduction of the spatial dependence of $w \in W$.

The essential ingredient of the proof of this paper may be summarized as follows:

- Discovery of the point process with last-arrival-time dependent intensity [11, 12]. The hydrodynamic limit turned out to have an expression in terms of probabilities of the process. This expression was absent in [14], which was a partial cause of

a technical boundedness assumptions on W . The process lack independent increment properties, which is a remnant of stochastic dependence among the particles through the position dependence of $w \in W$. This is to be compared with the earlier studies for position independent w , where the corresponding quantities have explicit formulas using exponentials of integration of w , related to the probabilities of the Poisson process [6, 5].

- Discovery of an intermediate model, defined in § 3, which we call the *flow driven stochastic ranking process*. The ‘intensity densities’ for the flow driven stochastic ranking process are involved but without position dependence, hence the distribution function φ for this model is a sum of independent processes, and a standard law of large numbers has a chance of explaining the fluctuation cancellation mechanism. (Introduction of an intermediate model resembles a notion of local equilibrium which appears in the hydrodynamic limit for diffusions. The limit of the stochastic ranking process is, in terms of fluid dynamics, a one-sided flow with evaporation from upper stream $y = 0$ to the down stream $y = 1$, and the correspondence to diffusion is only a kind of metaphor.)

Since the stochastic dependence among the particles induced by the move-to-front rule is handled by introduction of distribution functions φ , the real challenge from a viewpoint of mathematical analysis is the stochastic dependence through position dependence of $w \in W$, which was first studied in [14]. In the reference we adopted a sophisticated method (than adopted in this paper) based on submartingale inequalities, which worked well with the restricting assumptions in the reference, and enabled a relatively quick proof without introducing intermediate models.

- Development and application of a uniform strong complete law of large numbers for independent monotone function valued random variables.

Since the flow driven stochastic ranking process is stochastically similar to the stochastic ranking process with position independent intensities, a strong law of large numbers for independent processes is applicable. Since, however, we later need to compare this intermediate model with the original model, we apply uniform convergence results stronger than in earlier works [5], where the law of large numbers for the independent processes was practically the final goal.

- Application of a hierarchy of multi time Gronwall type inequalities. The last step is to prove that the original stochastic ranking process has a same limit with the (appropriately chosen) flow driven stochastic ranking process. To evaluate the difference, we couple the two models, and resort to a multi time variable recursive version of Gronwall type inequality which we develop in § 4.

In [14], we also proved the occurrence of propagation of chaos. Namely, for each integer L , the tagged particle system

$$(Y_1^{(N)}(t), Y_2^{(N)}(t), \dots, Y_L^{(N)}(t))$$

converges to a limit process uniformly in $t \in [0, T]$ as $N \rightarrow \infty$, if the system of the initial positions $(y_1^{(N)}, y_2^{(N)}, \dots, y_L^{(N)})$ converges, and the components of the limit are independent of each other. A corresponding result is also proved in this paper.

The plan of the present paper is as follows. In § 2 we state the main results, that the stochastic ranking process with space-time dependent intensities has a limit characterized by the point processes with last-arrival-time dependent intensities. For convenience, we will summarize the definition and relevant results of the point processes in § A. In § 3 we formulate the flow driven stochastic ranking process and prove the existence of the strong uniform law of large numbers (the large particle number limit). In § 4 we formulate and prove a hierarchy of a multi time version of the Gronwall's inequality, which we use in § 5 to complete a proof of the main theorem stated in § 2.

Acknowledgements.

The author would like to thank Professor Seiichiro Kusuoka, Professor Kazumasa Kuwada, and Professor Naoto Miyoshi, for their interest in the present work and for giving the author opportunities for seminar talks at Okayama University, University of Tokyo, and Tokyo Institute of Technology, respectively. The author in particular thanks Prof. Kusuoka for discussions and collaborations.

2 Formulation and the main results.

To state the main results precisely, we first formulate the quantities which appear in the infinite particle limit of the stochastic ranking process. Denote the sets of initial ($t = 0$) points in the space-time $[0, 1] \times [0, T]$, the set of upper stream boundary ($y = 0$) points, and their union, the set of initial/boundary points, respectively by

$$(9) \quad \begin{aligned} \Gamma_b &= \{(0, s) \mid 0 \leq s \leq T\}, \\ \Gamma_i &= \{(z, 0) \mid 0 \leq z \leq 1\}, \\ \Gamma &= \Gamma_b \cup \Gamma_i. \end{aligned}$$

For $t \in [0, T]$, denote the set of initial/boundary points up to time t by

$$(10) \quad \Gamma_t = \{(z, t_0) \in \Gamma \mid t_0 \leq t\} = \Gamma_i \cup \{(0, t_0) \in \Gamma_b \mid 0 \leq t_0 \leq t\},$$

and the set of admissible pairs of the initial/boundary point γ and time t by

$$(11) \quad \Delta_T = \{(\gamma, t) \in \Gamma_T \times [0, T] \mid \gamma \in \Gamma_t\}.$$

Define the set of *flows* Θ_T on $[0, 1] \times [0, T]$ by

$$(12) \quad \Theta_T = \{ \theta : \Delta_T \rightarrow [0, 1] \mid \text{continuous, } \theta((y_0, t_0), t_0) = y_0, (y_0, t_0) \in \Gamma_T, \\ \text{surjective and non-increasing in } \gamma \text{ for each } t, \\ \text{non-decreasing in } t \text{ for each } \gamma \},$$

where, we define a total order \succeq on the initial/boundary set Γ_T by

$$(13) \quad s \leq t, z \leq y \Leftrightarrow (0, T) \succeq (0, t) \succeq (0, s) \succeq (0, 0) \succeq (z, 0) \succeq (y, 0) \succeq (1, 0).$$

For example,

$$(14) \quad \theta((1, 0), t) = 1, \quad t \in [0, T], \quad \theta \in \Theta_T.$$

For each $\theta \in \Theta_T$, $w \in W$, and $z \in [0, 1]$, define $\tilde{w}_{\theta, w, z}$, a non-negative valued continuous function of (s, t) satisfying $0 \leq s \leq t \leq T$, by

$$(15) \quad \tilde{w}_{\theta, w, z}(s, t) = \begin{cases} w(\theta((z, 0), t), t), & \text{if } s = 0, \\ w(\theta((0, s), t), t), & \text{if } s > 0. \end{cases}$$

Note that $\tilde{w}_{\theta, w, z}$ is independent of z if $s > 0$. Let $\theta \in \Theta_T$, and put

$$(16) \quad \varphi_\theta(dw, \gamma, t) = \int_{z \in [y_0, 1]} \mathbb{P}[\tilde{v}_{\theta, w, z}(t) = \tilde{v}_{\theta, w, z}(t_0)] \mu_0(dw \times dz), \\ \gamma = (y_0, t_0) \in \Gamma_t, (\gamma, t) \in \Delta_T,$$

where $\tilde{v}_{\theta, w, z}$ is the point process with last-arrival-time dependent intensity, denoted by N in § A, with the ‘intensity density’ $\omega(s, t)$ in the definition (141) of the process given by $\tilde{w}_{\theta, w, z}(s, t)$ of (15), and μ_0 is a Borel probability measure on the direct product space $W \times [0, 1]$.

The following is proved in [11].

THEOREM 2.1 ([11, Theorem 9, (89)]). *There exists a unique flow $y_C \in \Theta_T$ such that*

$$(17) \quad \theta(\gamma, t) = 1 - \varphi_\theta(W, \gamma, t), \quad \gamma = (y_0, t_0) \in \Gamma, (\gamma, t) \in \Delta_T,$$

holds for $\theta = y_C$.

In [11] below the Theorem it is remarked that there exists μ_t , taking values in the space of Borel probability measures on $W \times [0, 1]$, such that

$$(18) \quad \varphi_{y_C}(dw, (y_0, t_0), t) = \mu_t(dw \times [y_C((y_0, t_0), t), 1]), \\ ((y_0, t_0), t) \in \Delta_T,$$

and below Theorem 1 in [11] it is also remarked that μ_t and y_C satisfy

$$(19) \quad y_C(\gamma, t) = y_0 + \int_{t_0}^t \int_{W \times [y_C(\gamma, s), 1]} w(z, s) \mu_s(dw \times dz) ds.$$

Next we state our assumptions on the infinite particle limit $N \rightarrow \infty$ of initial ($t = 0$) conditions. We will assume a standard supremum norm on the space of continuous functions on the closed interval $[0, 1] \times [0, T]$, with which we define the weak convergence of Borel probability measures on W in the standard way [2]. For the initial distribution $\mu_0^{(N)}$ in (4), we assume the following weak convergence with additional uniform bounds on the order of convergence to μ_0 in (16):

$$(20) \quad \begin{aligned} & \text{For any set } H \subset C^0(W \times [0, 1]; \mathbb{R}) \text{ of uniformly bounded,} \\ & \text{equicontinuous functions, } \exists \delta \in (0, \frac{1}{2}), \exists C > 0; \\ & (\forall N \in \mathbb{N}) (\forall \tilde{h} \in H) (\forall y \in [0, 1]) \\ & \left| \int_{W \times [y, 1]} \tilde{h}(w, z) \mu_0^{(N)}(dw \times dy) - \int_{W \times [y, 1]} \tilde{h}(w, z) \mu_0(dw \times dy) \right| \leq \frac{C}{N^\delta}. \end{aligned}$$

Denote the marginal distribution of μ_0 on W by λ ;

$$(21) \quad \lambda(dw) = \mu_0(dw \times [0, 1]),$$

and put

$$(22) \quad \lambda^{(N)} = \frac{1}{N} \sum_{i=1}^N \delta_{w_i}.$$

Comparing with (3), we see that $\lambda^{(N)}$ is the marginal distribution of $\mu_t^{(N)}$ on W for all t ;

$$(23) \quad \lambda^{(N)}(dw) = \mu_t^{(N)}(dw \times [0, 1]), \quad t \in [0, T].$$

For λ we assume

$$(24) \quad M_W := \int_W \|w\|_T \lambda(dw) < \infty,$$

where, and hereafter, we put

$$(25) \quad \|f\|_T = \sup_{(z, s) \in [0, 1] \times [0, T]} |f(z, s)|.$$

The assumption (20) implies, with (23) and (21),

$$(26) \quad \lambda^{(N)} \rightarrow \lambda, \text{ weakly as } N \rightarrow \infty.$$

In addition we assume convergence of the average of $\|w\|_T$:

$$(27) \quad \lim_{N \rightarrow \infty} \int_W \|w\|_T \lambda^{(N)}(dw) = M_W.$$

REMARK 2.2. In (20) we assume uniform order of convergence $O(N^{-\delta})$, while Ascoli–Arzelà type theorem implies uniform convergence but has no control in general on the order of convergence.

We are ready to state the main results of this paper.

THEOREM 2.3 (Main Theorem). *Under the assumptions (8), (20), (24), and (27), with probability 1, $\mu_t^{(N)} \rightarrow \mu_t$, weakly as $N \rightarrow \infty$, uniformly in t , where μ_t is as in (18). Explicitly, we prove*

$$(28) \quad \lim_{N \rightarrow \infty} \sup_{t \in [0, T]} \left| \int_W h(w) \mu_t^{(N)}(dw \times [y, 1]) - \int_W h(w) \mu_t(dw \times [y, 1]) \right| = 0, \text{ a.s.},$$

for all $y \in [0, 1]$ and bounded continuous function $h : W \rightarrow \mathbb{R}$.

Assume in addition that for a positive integer L and $y_i \in [0, 1)$, $i = 1, 2, \dots, L$,

$$(29) \quad \nu_i^{(N)} = \nu_i, \quad N \in \mathbb{N}, \quad \text{and} \quad \lim_{N \rightarrow \infty} y_i^{(N)} = y_i, \quad \text{for } i = 1, 2, \dots, L.$$

Then, with probability 1, the tagged particle system

$$(Y_1^{(N)}(t), Y_2^{(N)}(t), \dots, Y_L^{(N)}(t))$$

converges as $N \rightarrow \infty$, uniformly in $t \in [0, T]$ to a limit $(Y_1(t), Y_2(t), \dots, Y_L(t))$. Here, for each $i = 1, 2, \dots, L$, Y_i is the unique solution to

$$(30) \quad \begin{aligned} Y_i(t) = & y_i + \int_{s \in (0, t]} \int_{(w, z) \in W \times [Y_i(s-), 1]} w(z, s) \mu_s(dw \times dz) ds \\ & - \int_{s \in (0, t]} \int_{\xi \in [0, \infty)} Y_i(s-) \mathbf{1}_{\xi \in [0, w_i(Y_i(s-), s))} \nu_i(d\xi ds), \\ & i = 1, 2, \dots, L, \quad t \in [0, T]. \end{aligned}$$

Theorem 2.3 implies propagation of chaos for the stochastic ranking processes. For each N all of $\{Y_i^{(N)}\}$ are random and interact with each other and $\mu_t^{(N)}$ is also random. However, the limit μ_t is deterministic. Furthermore, the randomness of the limit process Y_i of a tagged particle depends only on its associated Poisson random measure ν_i , and is independent of Y_j or ν_j with $j \neq i$.

Incidentally, we proved almost sure convergence for $\mu_t^{(N)}$ in (28), while we made no assumptions on the relation between the set of measures $\{\nu_i^{(N)} \mid i = 1, 2, \dots, N\}$ for different N . This is stronger than the law of large numbers appearing in the context of random walks $\frac{1}{N} \sum_{i=1}^N X_i$, where each X_i is fixed for all N , while (28) corresponds to considering $\frac{1}{N} \sum_{i=1}^N X_i^{(N)}$, where we assume no relation among $X_i^{(N)}$ for different N . Such a type of convergence is known for sums of real valued random variables as complete convergence [15, 3, 4]. In this context, (18) is an example of complete convergence for a sequence of measure valued random variables.

3 Infinite particle limit of flow driven stochastic ranking process.

3.1 Flow driven stochastic ranking process.

In this section, we introduce an intermediate model which we use to prove convergence of $\mu_t^{(N)}$ in Theorem 2.3.

Let $\{w_i\}$, $\{y_i^{(N)}\}$, and $\{\nu_i^{(N)}\}$ be as in the stochastic ranking process (5). Let $\theta \in \Theta_T$ be a flow, and for each $i = 1, 2, \dots, N$, let $\tilde{\nu}_i^{(N,\theta)}$ be a point process of N in (147), with $\nu = \nu_i^{(N)}$ and $\omega = \tilde{w}_{\theta, w_i, y_i^{(N)}}$, where the last notation is as in (15) with $w = w_i$ and $z = y_i^{(N)}$. Define a system of stochastic processes $Y_i^{(N,\theta)}$, $i = 1, 2, \dots, N$, by

$$(31) \quad \begin{aligned} Y_i^{(N,\theta)}(t) &= y_i^{(N)} + \frac{1}{N} \sum_{j=1}^N \int_{s \in (0,t]} \mathbf{1}_{Y_j^{(N,\theta)}(s-) > Y_i^{(N,\theta)}(s-)} \tilde{\nu}_j^{(N,\theta)}(ds) \\ &\quad - \int_{s \in (0,t]} Y_i^{(N,\theta)}(s-) \tilde{\nu}_i^{(N,\theta)}(ds), \quad i = 1, 2, \dots, N, \quad t \geq 0. \end{aligned}$$

We will call this system, the stochastic ranking process driven by the flow θ .

The fluctuation of (31) is coupled to those of the stochastic ranking process of (1) via the set of Poisson random measures $\{\nu_i^{(N)}\}$. Using (144) and (147), we have an expression of $\tilde{\nu}_i^{(N,\theta)}$ using $\nu_i^{(N)}$. Define a sequence of stopping times, $0 = \tau_{i,0}^{(N,\theta)} < \tau_{i,1}^{(N,\theta)} < \dots$, by

$$(32) \quad \begin{aligned} \tau_{i,0}^{(N,\theta)} &= 0, \\ \tau_{i,k+1}^{(N,\theta)} &= \inf\{t > \tau_{i,k}^{(N,\theta)} \mid \nu_i^{(N)}(\{(s, \xi) \in (\tau_{i,k}^{(N,\theta)}, T] \times [0, \infty) \mid \\ &\quad 0 \leq \xi \leq w_i(Y_i^{(N,\theta)}(s-), s)\}) > 0\}, \quad k \in \mathbb{Z}_+. \end{aligned}$$

$\tau_{i,k}^{(N,\theta)}$ is the time that the particle i in the flow driven stochastic ranking process jumps to the top for the k -th time. Put

$$(33) \quad \gamma_i^{(N,\theta)}(t) = \begin{cases} (y_i^{(N)}, 0), & 0 \leq t < \tau_{i,1}^{(N,\theta)}, \\ (0, \tau_{i,k}^{(N,\theta)}), & \tau_{i,k}^{(N,\theta)} \leq t < \tau_{i,k+1}^{(N,\theta)}, \quad k = 1, 2, \dots \end{cases}$$

Using $\gamma_i^{(N,\theta)}(t)$, we have an expression

$$(34) \quad \tilde{\nu}_i^{(N,\theta)}(t) = \int_{s \in (0,t]} \int_{\xi \in [0, \infty)} \mathbf{1}_{\xi \in [0, w_i(\theta(\gamma_i^{(N,\theta)}(s-), s-), s))} \nu_i^{(N)}(d\xi ds),$$

and substituting (34) in (31) we further have

$$\begin{aligned}
(35) \quad Y_i^{(N,\theta)}(t) &= y_i^{(N)} \\
&+ \frac{1}{N} \sum_{j=1}^N \int_{s \in (0,t]} \int_{\xi \in [0,\infty)} \mathbf{1}_{Y_j^{(N,\theta)}(s-) > Y_i^{(N,\theta)}(s-)} \\
&\times \mathbf{1}_{\xi \in [0, w_j(\theta(\gamma_j^{(N,\theta)}(s-), s-), s))} \nu_j^{(N)}(d\xi ds) \\
&- \int_{s \in (0,t]} \int_{\xi \in [0,\infty)} Y_i^{(N,\theta)}(s-) \mathbf{1}_{\xi \in [0, w_i(\theta(\gamma_i^{(N,\theta)}(s-), s-), s))} \nu_i^{(N)}(d\xi ds), \\
&i = 1, 2, \dots, N, \quad t \geq 0.
\end{aligned}$$

This is to be compared with the (original) stochastic ranking process (1). We see that (35) is obtained from (1) by replacing $Y_j^{(N)}(s)$ appearing as a variable for w_j by $\theta(\gamma_j^{(N,\theta)}(s), s)$, and otherwise, by $Y_j^{(N,\theta)}(s)$.

3.2 Characteristic curve and distribution function.

For each $i = 1, 2, \dots, N$ and $0 \leq t_0 \leq t$, define an event $J_i^{(N,\theta)}(t_0, t) \subset \Omega$ by

$$(36) \quad J_i^{(N,\theta)}(t_0, t) = \{\omega \in \Omega \mid \tilde{\nu}_i^{(N,\theta)}(t)(\omega) > \tilde{\nu}_i^{(N,\theta)}(t_0)(\omega)\},$$

Since $\tilde{\nu}^{(N,\theta)}$ is increasing, the complement is

$$(37) \quad J_i^{(N,\theta)}(t_0, t)^c = \{\omega \in \Omega \mid \tilde{\nu}_i^{(N,\theta)}(t)(\omega) = \tilde{\nu}_i^{(N,\theta)}(t_0)(\omega)\}.$$

On the event $J_i^{(N,\theta)}(t_0, t)^c$, the contribution of the last term on the right hand side of (31) to the difference $Y_i^{(N,\theta)}(t) - Y_i^{(N,\theta)}(t_0)$ is 0, hence $Y_i^{(N,\theta)}$ is non-decreasing in the interval $[t_0, t]$. In other words, $J_i^{(N,\theta)}(t_0, t)$ of (36) is the event that the particle i jumps to the top $y = 0$ during $(t_0, t]$.

Define the characteristic curve for the stochastic ranking process driven by the flow θ by

$$(38) \quad Y_C^{(N,\theta)}((y_0, t_0), t) = y_0 + \frac{1}{N} \sum_{j \in [1, N]; Y_j^{(N,\theta)}(t_0) \geq y_0} \mathbf{1}_{J_j^{(N,\theta)}(t_0, t)},$$

for $(y_0, t_0) \in [0, 1] \times [0, T]$ and $t \geq t_0$. For example,

$$(39) \quad Y_C^{(N,\theta)}((1, t_0), t) = 1, \quad t \geq t_0 \geq 0.$$

Since $NY_j^{(N,\theta)}$ takes integer values, we can write (38) using (37) as

$$(40) \quad Y_C^{(N,\theta)}((y_0, t_0), t) = y_0 + \frac{[N(1 - y_0)]}{N} - \frac{1}{N} \sum_{j; Y_j^{(N,\theta)}(t_0) \geq y_0} \mathbf{1}_{J_j^{(N,\theta)}(t_0, t)^c},$$

where $[x]$ is the largest integer not exceeding x .

We note the following expression corresponding to (5).

LEMMA 3.1. (i) For $t \geq t_0 \geq 0$ and $0 \leq y_0 \leq 1$, it holds that

$$(41) \quad Y_C^{(N,\theta)}((y_0, t_0), t) = y_0 + \frac{1}{N} \sum_{j=1}^N \int_{s \in (t_0, t]} \int_{\xi \in [0, \infty)} \mathbf{1}_{Y_j^{(N,\theta)}(s-) \geq Y_C^{(N,\theta)}((y_0, t_0), s-)} \mathbf{1}_{\xi \in [0, w_j(\theta(\gamma_j^{(N,\theta)}(s-), s), s))} \nu_j^{(N)}(d\xi ds),$$

(ii) It holds that

$$(42) \quad Y_i^{(N,\theta)}(t) = Y_C^{(N,\theta)}(\gamma_i^{(N,\theta)}(t), t), \quad t \in [0, T],$$

where $\gamma_i^{(N,\theta)}$ is as in (33).

PROOF. Since on the event $J_i^{(N,\theta)}(t_0, t)^c$, the contribution of the last term on the right hand side of (31) to the difference $Y_i^{(N,\theta)}(t) - Y_i^{(N,\theta)}(t_0)$ disappears, $Y_i^{(N,\theta)}(t) - Y_i^{(N,\theta)}(t_0)$ and

$$Y_C^{(N,\theta)}((Y_i^{(N,\theta)}(t_0), t_0), t) - Y_i^{(N,\theta)}(t_0)$$

should satisfy the same equation for $[t_0, t]$ on $J_i^{(N,\theta)}(t_0, t)^c$. This implies (41).

By definition (33), $J_i^{(N,\theta)}(t_0, t)^c$ holds if $\gamma_i^{(N,\theta)}(t) = (y_0, t_0)$. Hence (42) follows. \square

In analogy to (3), for each positive integer N define a joint empirical distribution of jump rate and position of particles on $W \times [0, 1]$ by

$$(43) \quad \mu_t^{(N,\theta)} = \frac{1}{N} \sum_{i=1}^N \delta_{(w_i, Y_i^{(N,\theta)}(t))} \quad t \in [0, T].$$

When integrated over position, we recover the distribution $\lambda^{(N)}$ of jump rates for the (original) stochastic ranking process (22), independently of t and θ ;

$$(44) \quad \mu_t^{(N,\theta)}(dw \times [0, 1]) = \lambda^{(N)}(dw).$$

The initial distribution of $\mu_t^{(N,\theta)}$, found by substituting (31) at $t = 0$ to (43), coincide with that of the original model $\mu_0^{(N)}$ in (4):

$$(45) \quad \mu_0^{(N,\theta)} = \mu_0^{(N)}.$$

A set of spatial distribution functions $\varphi^{(N,\theta)}$ is defined analogously to (6), by a convolution of (38) and (43);

$$(46) \quad \begin{aligned} \varphi^{(N,\theta)}(dw, \gamma, t) &= \mu_t^{(N,\theta)}(dw \times [Y_C^{(N,\theta)}(\gamma, t), 1]), \\ \gamma = (y_0, t_0) &\in [0, 1] \times [0, T], \quad t \geq t_0. \end{aligned}$$

$\varphi^{(N,\theta)}(dw, \gamma, t)$ denotes the empirical distribution of jump rates (intensity densities) of those particles which was in a downstream of y_0 at time t_0 that have not jumped to the top in the time period $(t_0, t]$. For $t = t_0$ we have

$$(47) \quad \varphi^{(N,\theta)}(dw, (y_0, t_0), t_0) = \mu_{t_0}^{(N,\theta)}(dw \times [y_0, 1]), \quad (y_0, t_0) \in [0, 1] \times [0, T].$$

$\varphi^{(N,\theta)}$ is a refinement of $Y_C^{(N,\theta)}$ in the following sense. First, (46) and (43) imply

$$\varphi^{(N,\theta)}(dw, \gamma, t) = \frac{1}{N} \sum_{j=1}^N \mathbf{1}_{Y_j^{(N,\theta)}(t) \geq Y_C^{(N,\theta)}(\gamma, t)} \delta_{w_j}(dw).$$

Next, $Y_j^{(N,\theta)}(t) \geq Y_C^{(N,\theta)}(\gamma, t)$ occurs if and only if the inequality holds at $t = t_0$ and j does not jump to the top in the interval $(t_0, t]$. Hence (36) implies

$$(48) \quad \varphi^{(N,\theta)}(dw, (y_0, t_0), t) = \frac{1}{N} \sum_{j; Y_j^{(N,\theta)}(t_0) \geq y_0} \mathbf{1}_{J_j^{(N,\theta)}(t_0, t)^c} \delta_{w_j}(dw).$$

Comparing (48) with (40), we have

$$(49) \quad \begin{aligned} Y_C^{(N,\theta)}(\gamma, t) &= y_0 + \frac{[N(1 - y_0)]}{N} - \varphi^{(N,\theta)}(W, \gamma, t), \\ \gamma &= (y_0, t_0) \in [0, 1] \times [0, T], \quad t \geq t_0. \end{aligned}$$

The spatial distribution function $\varphi^{(N,\theta)}(dw, \gamma, t)$ is defined for any $\gamma = (y_0, t_0) \in [0, 1] \times [0, T]$ satisfying $t \geq t_0$, but is particularly important when $\gamma \in \Gamma_t$. In fact, both for the case $\gamma = (y_0, 0) \in \Gamma_i$ and the case $\gamma = (0, t_0) \in \Gamma_t$ we have

$$(50) \quad \begin{aligned} \varphi^{(N,\theta)}(dw, (y_0, t_0), t) &= \frac{1}{N} \sum_{j; y_j^{(N)} \geq y_0} \mathbf{1}_{J_j^{(N,\theta)}(t_0, t)^c} \delta_{w_j}(dw), \\ \gamma &= (y_0, t_0) \in \Gamma_t, \quad t \in [0, T]. \end{aligned}$$

Note the conditions of the summation, which is non-random in (50), while is random in (48). Since $J_j^{(N,\theta)}(t_0, t)$ is independent of $\nu_i^{(N)}$, $i \neq j$, (50) implies that $\varphi^{(N,\theta)}(dw, \gamma, t)$ is an arithmetic average of independent random variables, if $\gamma \in \Gamma_t$. With this fact, we restrict the domain of definition to Δ_T of (11) in § 3.3.

3.3 Convergence of distribution function.

3.3.1 Statement of the theorem.

Here we will state a strong law of large numbers for the spatial distribution function $\varphi^{(N,\theta)}$ of the flow driven stochastic ranking process, uniform in initial point γ and time t .

For a bounded continuous function $h : W \rightarrow \mathbb{R}$, we put

$$(51) \quad C_h = \sup_{w \in W} |h(w)| < \infty,$$

and use a notation

$$(52) \quad \varphi^{(N,\theta)}(h, \gamma, t) = \int_W h(w) \varphi^{(N,\theta)}(dw, \gamma, t).$$

With (48) we have

$$(53) \quad \varphi^{(N,\theta)}(h, \gamma, t) = \frac{1}{N} \sum_{j; Y_j^{(N,\theta)}(t_0) \geq y_0} h(w_j) \mathbf{1}_{J_j^{(N,\theta)}(t_0, t)^c}, \quad \gamma = (y_0, t_0).$$

As in (52) we also use a notation

$$(54) \quad \varphi_\theta(h, \gamma, t) = \int_W h(w) \varphi_\theta(dw, \gamma, t).$$

for φ_θ in (16).

THEOREM 3.2. *Assume (8), (20), (24), and (27). Then for any $p > 0$ there exists a positive constant C depending only on p and δ , (and is independent of N , θ , and h .) such that for any bounded continuous $h : W \rightarrow \mathbb{R}$,*

$$(55) \quad \mathbb{E} \left[\sup_{(\gamma, t) \in \Delta_T} \left| \varphi^{(N,\theta)}(h, \gamma, t) - \varphi_\theta(h, \gamma, t) \right|^{2p} \right] \leq \frac{C C_h^{2p}}{N^{2p\delta}}, \quad N \in \mathbb{N},$$

holds, where C_h is as in (51).

Theorem 2.1, Theorem 3.2 with $h = \mathbf{1}_W$ and $\theta = y_C$, and (49) imply the following.

COROLLARY 3.3. *Under the assumptions of Theorem 3.2, for any $p > 0$ there exists a positive constant C depending only on p and δ , such that*

$$(56) \quad \mathbb{E} \left[\sup_{(\gamma, t) \in \Delta_T} \left| Y_C^{(N, y_C)}(\gamma, t) - y_C(\gamma, t) \right|^{2p} \right] \leq \frac{C}{N^{2p\delta}}, \quad N \in \mathbb{N}.$$

Among $\theta \in \Theta$, $\theta = y_C$ is the only flow that satisfies (56).

We can take $p > \frac{1}{2\delta}$ in Theorem 3.2, from which we have

$$\mathbb{E} \left[\sum_{N=1}^{\infty} \sup_{(\gamma, t) \in \Delta_T} \left| \varphi^{(N,\theta)}(h, \gamma, t) - \varphi_\theta(h, \gamma, t) \right|^{2p} \right] \leq C \sum_{N=1}^{\infty} \frac{1}{N^{2p\delta}} < \infty,$$

which implies

$$\sum_{N=1}^{\infty} \sup_{(\gamma, t) \in \Delta_T} \left| \varphi^{(N,\theta)}(h, \gamma, t) - \varphi_\theta(h, \gamma, t) \right|^{2p} < \infty$$

for almost all $\omega \in \Omega$. For such ω we then have (by continuity and increasing property of $x \mapsto x^{2p}$ for $x \geq 0$),

$$\begin{aligned} & \lim_{N \rightarrow \infty} \sup_{(\gamma, t) \in \Delta_T} \left| \varphi^{(N, \theta)}(h, \gamma, t) - \varphi_\theta(h, \gamma, t) \right| \\ &= \left(\lim_{N \rightarrow \infty} \sup_{(\gamma, t) \in \Delta_T} \left| \varphi^{(N, \theta)}(h, \gamma, t) - \varphi_\theta(h, \gamma, t) \right|^{2p} \right)^{1/(2p)} \\ &= 0, \end{aligned}$$

hence

$$\lim_{N \rightarrow \infty} \sup_{(\gamma, t) \in \Delta_T} \left| \varphi^{(N, \theta)}(h, \gamma, t) - \varphi_\theta(h, \gamma, t) \right| = 0, \quad a.e..$$

This implies convergence of joint empirical distribution $\mu_t^{(N, \theta)}$ to $\mu_{y_C, t}$, a deterministic distribution determined by $\varphi_{y_C}(h, \gamma, t)$. Similarly, we have convergence of characteristic curves from Corollary 3.3.

For $\theta \neq y_C$, the stochastic ranking process driven by the flow θ converges to a limit with deterministic distribution, but the resulting trajectories of particles are different from the given flow θ , due to the uniqueness result of Theorem 2.1. Only when $\theta = y_C$ we have the limit trajectories equal to the given flow y_C .

A proof of Theorem 3.2 is composed of 2 parts. In § 3.3.2 we prove that $\varphi^{(N, \theta)} - \mathbb{E}[\varphi^{(N, \theta)}]$ converges to 0, using a strong uniform law of large numbers in [13]. In § 3.3.3 we prove that $\mathbb{E}[\varphi^{(N, \theta)}]$ converges to φ_θ , using the estimates in [11]. Relevant results of [11] are summarized in § A for convenience.

3.3.2 Strong uniform law of large numbers.

Here we will prove the following.

PROPOSITION 3.4. *Assume (8), (20), (24), and (27). Then for any p and δ satisfying*

$$(57) \quad p > 0, \quad 0 < \delta < \frac{1}{2},$$

there exists a positive constant C (independent of N , θ , and h ,) such that for any bounded continuous $h : W \rightarrow \mathbb{R}$

$$(58) \quad \mathbb{E} \left[\sup_{(\gamma, t) \in \Delta_T} \left| \varphi^{(N, \theta)}(h, \gamma, t) - \mathbb{E}[\varphi^{(N, \theta)}(h, \gamma, t)] \right|^{2p} \right] \leq \frac{C C_h^{2p}}{N^{2p\delta}}, \quad N \in \mathbb{N},$$

where C_h is as in (51).

PROOF. Put $h_{\pm} = (\pm h) \vee 0$, so that $h = h_+ - h_-$ decomposes the function h to positive and negative parts. Applying (48), (11), (10), and triangular inequality in the form

$$(a + b)^q \leq (2^{q-1} \vee 1)(a^q + b^q), \quad a \geq 0, \quad b \geq 0, \quad q > 0,$$

we have, with $\gamma = (y_0, t_0) \in \Gamma_t$,

$$\begin{aligned}
& \mathbb{E} \left[\sup_{((y_0, t_0), t) \in \Delta_T} \left| \varphi^{(N, \theta)}(h, \gamma, t) - \mathbb{E} \left[\varphi^{(N, \theta)}(h, \gamma, t) \right] \right|^q \right] \\
&= \mathbb{E} \left[\sup_{((y_0, t_0), t)} \left| \frac{1}{N} \sum_{Y_j^{(N, \theta)}(t_0) \geq y_0} (h_+(w_j) - h_-(w_j)) \right. \right. \\
&\quad \left. \left. \times (\mathbf{1}_{J_j^{(N, \theta)}(t_0, t)^c} - \mathbb{E}[\mathbf{1}_{J_j^{(N, \theta)}(t_0, t)^c}]) \right|^q \right] \\
(59) \quad &\leq (2^{q-1} \vee 1) \mathbb{E} \left[\sup_{((y_0, t_0), t)} \left| \frac{1}{N} \sum_{Y_j^{(N, \theta)}(t_0) \geq y_0} h_+(w_j) \right. \right. \\
&\quad \left. \left. \times (\mathbf{1}_{J_j^{(N, \theta)}(t_0, t)^c} - \mathbb{E}[\mathbf{1}_{J_j^{(N, \theta)}(t_0, t)^c}]) \right|^q \right] \\
&\quad + (2^{q-1} \vee 1) \mathbb{E} \left[\sup_{((y_0, t_0), t)} \left| \frac{1}{N} \sum_{Y_j^{(N, \theta)}(t_0) \geq y_0} h_-(w_j) \right. \right. \\
&\quad \left. \left. \times (\mathbf{1}_{J_j^{(N, \theta)}(t_0, t)^c} - \mathbb{E}[\mathbf{1}_{J_j^{(N, \theta)}(t_0, t)^c}]) \right|^q \right] \\
&\leq (2^{q-1} \vee 1) (R_{q,1,+}^{(N)} + R_{q,1,-}^{(N)} + R_{q,2,+}^{(N)} + R_{q,2,-}^{(N)}),
\end{aligned}$$

where

$$(60) \quad R_{q,1,\pm}^{(N)} = \mathbb{E} \left[\sup_{t \in [0, T]} \sup_{0 \leq y_0 < 1} \left| \frac{1}{N} \sum_{i: y_i^{(N)} \geq y_0} h_{\pm}(w_i) (\mathbf{1}_{J_i^{(N, \theta)}(0, t)^c} - \mathbb{E}[\mathbf{1}_{J_i^{(N, \theta)}(0, t)^c}]) \right|^q \right]$$

and

$$(61) \quad R_{q,2,\pm}^{(N)} = \mathbb{E} \left[\sup_{t \in [0, T]} \sup_{0 \leq t_0 < t} \left| \frac{1}{N} \sum_{i=1}^N h_{\pm}(w_i) (\mathbf{1}_{J_i^{(N, \theta)}(t_0, t)^c} - \mathbb{E}[\mathbf{1}_{J_i^{(N, \theta)}(t_0, t)^c}]) \right|^q \right].$$

To bound (60) and (61), we refer to the last theorem in [13, §2]. We reproduce the theorem in a specific form of

$$Z_i^{(N)}(s, t) = a_i^{(N)} \mathbf{1}_{\nu_i^{(N)}(t) > \nu_i^{(N)}(s)}$$

in place of $Z_i^{(N)}(s, t)$ in the reference.

PROPOSITION 3.5 ([13, Theorem 7]). *Let $T > 0$, and for each $N \in \mathbb{N}$, let $\nu_i^{(N)}$, $i = 1, \dots, N$, be a sequence of independent random variables taking values in a space of*

non-negative valued non-decreasing right continuous functions on $[0, T]$ with left limit. Let $r > 0$, and for $N \in \mathbb{N}$, let $M^{(N)} > 0$ and let $a_i^{(N)}$ and $w_i^{(N)}$, $i = 1, \dots, N$, be non-negative sequences. Assume that $a_i^{(N)} \leq M^{(N)}$, $i = 1, \dots, N$, and that

$$\begin{aligned} & |\mathbb{P}[\nu_i^{(N)}(t_2) > \nu_i^{(N)}(t_1)] - \mathbb{P}[\nu_i^{(N)}(s_2) > \nu_i^{(N)}(s_1)]| \\ & \leq w_i^{(N)}(|t_1 - s_1|^r + |t_2 - s_2|^r), \end{aligned}$$

for $0 \leq t_1 \leq t_2 \leq T$ and $0 \leq s_1 \leq s_2 \leq T$. Then for any $\delta \in (0, \frac{1}{2})$ and $p > 0$,

$$\begin{aligned} (62) \quad & \mathbb{E} \left[\sup_{t_2 \in [0, T]} \sup_{0 \leq t_1 < t_2} \left| \frac{1}{N} \sum_{i=1}^N a_i^{(N)} (\mathbf{1}_{\nu_i^{(N)}(t_2) > \nu_i^{(N)}(t_1)} - \mathbb{P}[\nu_i^{(N)}(t_2) > \nu_i^{(N)}(t_1)]) \right|^p \right]^{1/p} \\ & \leq \frac{M^{(N)}}{N^\delta} 2^{1-1/q} (C_q^q (2T(\bar{w}^{(N)})^{1/r} + 1) + 2^{2q})^{1/q}, \\ & N = N_0, N_0 + 1, \dots \end{aligned}$$

Here $q = q(p, \delta) = 3 \vee \frac{r+1}{r} \frac{2\delta}{1-2\delta} \vee p$, N_0 is the smallest integer satisfying $N_0^{rq/(2rq+2r+2)} \geq 2$, $\bar{w}^{(N)} = \frac{1}{N} \sum_{i=1}^N w_i^{(N)}$, and $C_q = \left(\frac{1}{2} (4k)^q + \frac{2k}{2k-q} (8k)^q \right)^{1/q}$ with k the smallest integer greater than $\frac{1}{2}q$, (In particular, q and N_0 are independent of N , $M^{(N)}$, and $\bar{w}^{(N)}$.)

To bound $R_{q,2,\pm}^{(N)}$ in (61), we apply Proposition 3.5 with $\nu_i^{(N)} = \tilde{\nu}_i^{(N,\theta)}$ and $a_i^{(N)} = h_\pm(w_i)$. Note that (152) and (15) with $w = w_i$ and $z = y_i^{(N)}$, and (25) imply

$$\begin{aligned} (63) \quad & 0 \leq -\frac{\partial}{\partial t} \mathbb{P}[\tilde{\nu}_i^{(N,\theta)}(t) = \tilde{\nu}_i^{(N,\theta)}(s)] \leq \left\| \tilde{w}_{\theta, w_i, y_i^{(N)}} \right\| \leq \|w_i\|_{\mathbb{T}}, \\ & 0 \leq \frac{\partial}{\partial s} \mathbb{P}[\tilde{\nu}_i^{(N,\theta)}(t) = \tilde{\nu}_i^{(N,\theta)}(s)] \leq \left\| \tilde{w}_{\theta, w_i, y_i^{(N)}} \right\| \leq \|w_i\|_{\mathbb{T}}. \end{aligned}$$

Recall (36) and (37). Comparing the left hand side of (62) with the right hand side of (61) with $q = 2p$, we see that we can apply Proposition 3.5 to $R_{2p,2,\pm}^{(N)}$ with

$$(64) \quad M^{(N)} = C_h, \quad r = 1, \quad w_i^{(N)} = \|w_i\|_{\mathbb{T}}.$$

Proposition 3.5 then implies that for any $\delta \in (0, \frac{1}{2})$ and $p > 0$,

$$(65) \quad R_{2p,2,\pm}^{(N)} \leq \frac{C_h^{2p}}{N^{2p\delta}} 2^{2p-1} \left(C_q^q (2T\bar{w}^{(N)} + 1) + 2^{2q} \right)^{2p/q}, \quad N > 2^{2+(4/q)},$$

where

$$\begin{aligned} q &= q(p, \delta) = 3 \vee \frac{4\delta}{1-2\delta} \vee (2p), \\ \bar{w}^{(N)} &= \frac{1}{N} \sum_{i=1}^N \|w_i\|_{\mathbb{T}} = \int_{\mathcal{W}} \|w\|_{\mathbb{T}} \lambda^{(N)}(dw), \end{aligned}$$

and C_q is a positive constant depending only on q .

Combining (64), (65), and (27), we see that there exists a positive constant $C_{p,\delta}$ independent of N , θ , and h , such that

$$(66) \quad R_{2p,2,\pm}^{(N)} \leq \frac{C_{p,\delta} C_h^{2p}}{N^{2p\delta}}, \quad N \in \mathbb{N}.$$

To bound $R_{q,1,\pm}^{(N)}$ in (60), we first note that

$$(67) \quad \begin{aligned} & R_{q,1,\pm}^{(N)} \\ & \leq \sum_{j=1}^N \mathbb{E} \left[\sup_{t \in [0,T]} \left| \frac{1}{N} \sum_{i; N y_i^{(N)} \geq j-1} h_{\pm}(w_i) (\mathbf{1}_{J_i^{(N,\theta)}(0,t)^c} - \mathbb{E}[\mathbf{1}_{J_i^{(N,\theta)}(0,t)^c}]) \right|^q \right] \\ & \leq \sum_{j=1}^N \mathbb{E} \left[\sup_{t \in [0,T]} \sup_{0 \leq t_0 < t} \left| \frac{1}{N} \sum_{i; N y_i^{(N)} \geq j-1} h_{\pm}(w_i) (\mathbf{1}_{J_i^{(N,\theta)}(t_0,t)^c} - \mathbb{E}[\mathbf{1}_{J_i^{(N,\theta)}(t_0,t)^c}]) \right|^q \right]. \end{aligned}$$

Comparing (67) with (61), we see that we can apply Proposition 3.5 with $\nu_i^{(N)} = \tilde{\nu}_i^{(N,\theta)}$ and

$$a_i^{(N)} = h_{\pm}(w_i) \mathbf{1}_{N y_i^{(N)} \geq j-1},$$

and (64), to the j -th term of the summation in the right hand side of (67), in a similar way as we did to $R_{2p,2,\pm}^{(N)}$. Using monotonicity of L^p norms with respect to p before applying Proposition 3.5, we have, for $q_0 \geq 2p$ and $\delta' \in (0, \frac{1}{2})$,

$$R_{2p,1,\pm} \leq R_{q_0,1,\pm}^{2p/q_0} \leq \frac{(C_{q_0/2,\delta'} C_h^{q_0})^{2p/q_0}}{N^{2p\delta' - 2p/q_0}}, \quad N \in \mathbb{N},$$

in place of (66). (The extra factor N compared to (66) is from the summation with respect to j in (67).) Now choose δ' and q_0 to satisfy $\delta < \delta' < \frac{1}{2}$ and $0 < \frac{1}{q_0} < \delta' - \delta$ to find

$$(68) \quad R_{2p,1,\pm} \leq R_{q_0,1,\pm}^{2p/q_0} \leq \frac{(C_{q_0/2,\delta'} C_h^{q_0})^{2p/q_0}}{N^{2p\delta}}, \quad N \in \mathbb{N}.$$

Proposition 3.4 finally follows from (59), (68), and (66). \square

3.3.3 Convergence of expectation.

Here we complete a proof of Theorem 3.2.

LEMMA 3.6. *Assume (27). Then, if $h : W \rightarrow \mathbb{R}$ is bounded and continuous,*

$$(69) \quad \lim_{N \rightarrow \infty} \int_W h(w) \|w\|_{\mathbb{T}} \lambda^{(N)}(dw) = \int_W h(w) \|w\|_{\mathbb{T}} \lambda(dw).$$

PROOF. Note first that for $M > 0$

$$(70) \quad a = (a \wedge M) + (a - M)_+$$

holds. Since $w \mapsto \|w\|_{\mathbb{T}} \wedge M$ is bounded and continuous, (26) (weak convergence) implies

$$(71) \quad \begin{aligned} \lim_{N \rightarrow \infty} \int_{\mathbb{W}} h(w) (\|w\|_{\mathbb{T}} \wedge M) \lambda^{(N)}(dw) &= \int_{\mathbb{W}} h(w) (\|w\|_{\mathbb{T}} \wedge M) \lambda(dw), \\ \lim_{N \rightarrow \infty} \int_{\mathbb{W}} \|w\|_{\mathbb{T}} \wedge M \lambda^{(N)}(dw) &= \int_{\mathbb{W}} \|w\|_{\mathbb{T}} \wedge M \lambda(dw), \end{aligned}$$

which, with convergence of expectation in (27), further implies

$$(72) \quad \lim_{N \rightarrow \infty} \int_{\mathbb{W}} (\|w\|_{\mathbb{T}} - M)_+ \lambda^{(N)}(dw) = \int_{\mathbb{W}} (\|w\|_{\mathbb{T}} - M)_+ \lambda(dw).$$

On the other hand, dominated convergence theorem, $0 \leq (\|w\|_{\mathbb{T}} - M)_+ \leq \|w\|_{\mathbb{T}}$, and (24) imply

$$\lim_{M \rightarrow \infty} \int_{\mathbb{W}} (\|w\|_{\mathbb{T}} - M)_+ \lambda(dw) = 0.$$

Hence, for any $\epsilon > 0$ there exists $M > 0$ such that $\int_{\mathbb{W}} (\|w\|_{\mathbb{T}} - M)_+ \lambda(dw) < \epsilon$. This and (72) further imply that there exists $N_0 > 0$ such that

$$\int_{\mathbb{W}} (\|w\|_{\mathbb{T}} - M)_+ \lambda^{(N)}(dw) < 2\epsilon, \quad N \geq N_0.$$

Put $C = \sup_{w \in \mathbb{W}} |h(w)| < \infty$. Then

$$\begin{aligned} & \left| \int_{\mathbb{W}} h(w) \|w\|_{\mathbb{T}} \lambda^{(N)}(dw) - \int_{\mathbb{W}} h(w) \|w\|_{\mathbb{T}} \lambda(dw) \right| \\ & \leq \left| \int_{\mathbb{W}} h(w) (\|w\|_{\mathbb{T}} \wedge M) \lambda^{(N)}(dw) - \int_{\mathbb{W}} h(w) (\|w\|_{\mathbb{T}} \wedge M) \lambda(dw) \right| \\ & \quad + C \int_{\mathbb{W}} (\|w\|_{\mathbb{T}} - M)_+ \lambda^{(N)}(dw) + C \int_{\mathbb{W}} (\|w\|_{\mathbb{T}} - M)_+ \lambda(dw) \\ & \leq \left| \int_{\mathbb{W}} h(w) (\|w\|_{\mathbb{T}} \wedge M) \lambda^{(N)}(dw) - \int_{\mathbb{W}} h(w) (\|w\|_{\mathbb{T}} \wedge M) \lambda(dw) \right| + 3C\epsilon. \end{aligned}$$

This and (71) imply

$$\overline{\lim}_{N \rightarrow \infty} \left| \int_{\mathbb{W}} h(w) \|w\|_{\mathbb{T}} \lambda^{(N)}(dw) - \int_{\mathbb{W}} h(w) \|w\|_{\mathbb{T}} \lambda(dw) \right| \leq 3C\epsilon.$$

Since the left hand side is independent of N , M , and ϵ , it must be 0. \square

Proof of Theorem 3.2. Let $h : W \rightarrow \mathbb{R}$ be a bounded continuous function, and put $C_h = \sup_{w \in W} |h(w)|$ as in (51). Proposition 3.4 implies that to prove Theorem 3.2, it suffices to prove

$$(73) \quad \sup_{(\gamma, t) \in \Delta_T} \left| \mathbb{E}[\varphi^{(N, \theta)}(h, \gamma, t)] - \varphi_\theta(h, \gamma, t) \right| \leq \frac{C C_h}{N^\delta}, \quad N \in \mathbb{N},$$

for C depending only on p and δ (independent of N , θ , and h).

Comparing the definition of $\tilde{v}_i^{(N, \theta)}$ given above (31) with that of $\tilde{v}_{\theta, w, z}$ given below (16), we see that $\tilde{v}_i^{(N, \theta)}$ and $\tilde{v}_{\theta, w_i, y_i^{(N)}}$ have identical distribution. Therefore, using (37) and (4) in (53), we have, for $t \geq t_0$,

$$(74) \quad \mathbb{E}[\varphi^{(N, \theta)}(h, (y_0, t_0), t)] = \int_{W \times [y_0, 1]} h(w) \mathbb{P}[\tilde{v}_{\theta, w, z}(t) = \tilde{v}_{\theta, w, z}(t_0)] \mu_0^{(N)}(dw \times dz).$$

Let $\gamma = (y_0, t_0) \in \Gamma$ and $(\gamma, t) \in \Delta_T$. (74) and (16) imply

$$\begin{aligned} & \mathbb{E}[\varphi^{(N, \theta)}(h, \gamma, t)] - \varphi_\theta(h, \gamma, t) \\ &= \int_{W \times [y_0, 1]} h(w) \mathbb{P}[\tilde{v}_{\theta, w, z}(t) = \tilde{v}_{\theta, w, z}(t_0)] (\mu_0^{(N)}(dw \times dz) - \mu_0(dw \times dz)). \end{aligned}$$

Hence,

$$(75) \quad \begin{aligned} & \sup_{(\gamma, t) \in \Delta_T} \left| \mathbb{E}[\varphi^{(N, \theta)}(h, \gamma, t)] - \varphi_\theta(h, \gamma, t) \right| \\ & \leq \sup_{((y_0, t_0), t) \in \Delta_T} \left| \int_{W \times [y_0, 1]} \tilde{h}_{t_0, t}(w, z) (\mu_0^{(N)}(dw \times dz) - \mu_0(dw \times dz)) \right|, \end{aligned}$$

where

$$(76) \quad \tilde{h}_{t_0, t}(w, z) = h(w) \mathbb{P}[\tilde{v}_{\theta, w, z}(t) = \tilde{v}_{\theta, w, z}(t_0)].$$

Choose the set H in (20) as the set of the functions $\tilde{h}_{t_0, t}$ in (76):

$$(77) \quad H = \{\tilde{h}_{t_0, t} : W \times [0, 1] \rightarrow \mathbb{R} \mid 0 \leq t_0 \leq t \leq T\}.$$

Uniform boundedness of the functions in H is obvious. If we prove that H is also equicontinuous, then by the assumption of Theorem 3.2 the consequence of (20) holds, which implies

$$\begin{aligned} & \sup_{y_0 \in [0, 1]} \sup_{0 \leq t_0 \leq t \leq T} \left| \int_{W \times [y_0, 1]} \tilde{h}_{t_0, t}(w, z) (\mu_0^{(N)}(dw \times dz) - \mu_0(dw \times dz)) \right| \leq \frac{C C_h}{N^\delta}, \\ & N \in \mathbb{N}. \end{aligned}$$

Applying this estimate to (75), we have (73), which proves Theorem 3.2.

We are left with proving equicontinuity of H .

First, for $(w, z) \in W \times [0, 1]$ and $\tilde{w}_{\theta, w, z}$ as in (15) (i.e., the ‘intensity density’ for $\tilde{\nu}_{\theta, w, z}$), and $0 \leq s \leq t \leq T$, put

$$(78) \quad \Omega_{\theta, w, z}(s, t) = \int_s^t \tilde{w}_{\theta, w, z}(s, u) du \quad \text{and} \quad \tilde{\Omega}_w(s, t) = \int_s^t w(1, u) du.$$

Then (8) and a mean value theorem imply

$$(79) \quad w(1, t) - C_W \leq \tilde{w}_{\theta, w, z}(s, t) \leq w(1, t) + C_W,$$

and

$$(80) \quad e^{-\tilde{\Omega}_w(s, t) - C_W(t-s)} \leq e^{-\Omega_{\theta, w, z}(s, t)} \leq e^{-\tilde{\Omega}_w(s, t) + C_W(t-s)}.$$

Note also an elementary formula in [11, (53)]

$$(81) \quad \int_{0 \leq u_1 \leq u_2 \leq \dots \leq u_k \leq s} \prod_{i=1}^k f(u_i) du_1 du_2 \dots du_k = \frac{1}{k!} \left(\int_0^s f(v) dv \right)^k,$$

valid for any integrable function $f : \mathbb{R} \rightarrow \mathbb{R}$, $s \geq 0$, and $k = 1, 2, \dots$,

A proof of equicontinuity of H now goes in a similar way as that of [11, Lemma 12]. Applying (151) to (76), we have

$$(82) \quad \begin{aligned} \tilde{h}_{t_0, t}(w, z) &= h(w) \sum_{k \geq 0} \int_{0 =: u_k < u_{k-1} < u_{k-2} < \dots < u_1 < u_0 \leq t_0} \\ &\times e^{-\sum_{i=0}^{k-1} \Omega_{\theta, w, z}(u_{i+1}, u_i) - \Omega_{\theta, w, z}(u_0, t)} \left(\prod_{i=0}^{k-1} \tilde{w}_{\theta, w, z}(u_{i+1}, u_i) du_i \right). \end{aligned}$$

Using (79), (80), and (78) to (82), while noting that (15) implies that $\tilde{w}_{\theta, w, z}(s, t)$ and $\Omega_{\theta, w, z}(s, t)$ is independent of z if $s > 0$, we have

$$|\tilde{h}_{t_0, t}(w, z') - \tilde{h}_{t_0, t}(w, z)| \leq I_{11}(z, z') + I_{12}(z, z'),$$

where

$$\begin{aligned} I_{11}(z, z') &= C_h e^{-\tilde{\Omega}_w(0, t) + C_W t} \sum_{k \geq 1} \int_{0 =: u_k < u_{k-1} < u_{k-2} < \dots < u_1 < u_0 \leq t_0} \\ &\times \left(\prod_{i=0}^{k-2} (w(1, u_i) + C_W) du_i \right) \\ &\times |\tilde{w}_{\theta, w, z'}(0, u_{k-1}) - \tilde{w}_{\theta, w, z}(0, u_{k-1})| du_{k-1}, \end{aligned}$$

and

$$\begin{aligned} I_{12}(z, z') &= C_h \sum_{k \geq 0} \int_{0 =: u_k < u_{k-1} < u_{k-2} < \dots < u_1 < u_0 \leq t_0} e^{-\tilde{\Omega}_w(u_{k-1}, t) + C_W(t - u_{k-1})} \\ &\times \left(\prod_{i=0}^{k-1} (w(1, u_i) + C_W) du_i \right) \\ &\times |e^{-\Omega_{\theta, w, z'}(0, u_{k-1})} - e^{-\Omega_{\theta, w, z}(0, u_{k-1})}| du_{k-1}. \end{aligned}$$

Using (15), (8), (81), and (78), we have

$$\begin{aligned} I_{11}(z, z') &\leq C_h C_W e^{-\tilde{\Omega}_w(t_0, t) + C_W(t+t_0)} \int_0^{t_0} |\theta((z', 0), v) - \theta((z, 0), v)| dv \\ &\leq C_h C_W e^{2C_W T} \int_0^T |\theta((z', 0), v) - \theta((z, 0), v)| dv. \end{aligned}$$

Using in addition

$$(83) \quad \begin{aligned} |e^{-x'} - e^{-x}| &= e^{-(x' \wedge x)} - e^{-(x' \vee x)} = e^{-(x' \wedge x)} (1 - e^{-|x' - x|}) \\ &\leq e^{-(x' \wedge x)} |x' - x| \leq e^{-x} e^{|x' - x|} |x' - x|, \end{aligned}$$

which follows from $|x' - x| = (x' \vee x) - (x' \wedge x) \geq x - (x \wedge x')$, we similarly have

$$\begin{aligned} I_{12}(z, z') &\leq C_h C_W e^{-\tilde{\Omega}_w(t_0, t) + C_W(t+t_0)} \int_0^{t_0} |\theta((z', 0), v) - \theta((z, 0), v)| dv \\ &\quad \times e^{C_W \int_0^{t_0} |\theta((z', 0), v) - \theta((z, 0), v)| dv} \\ &\leq C_h C_W e^{2C_W T} \int_0^T |\theta((z', 0), v) - \theta((z, 0), v)| dv \\ &\quad \times e^{C_W \int_0^T |\theta((z', 0), v) - \theta((z, 0), v)| dv}. \end{aligned}$$

Since the right hand sides of the bounds for I_{11} and I_{12} are uniform in t_0 and t , these prove equicontinuity in the variable $z \in [0, 1]$ of functions $\tilde{h}_{t_0, t}(w, z)$ in H .

In a similar way as the proof of equicontinuity with respect to z , we have

$$|\tilde{h}_{t_0, t}(w', z) - \tilde{h}_{t_0, t}(w, z)| \leq I_{21}(w, w') + I_{22}(w, w'),$$

where

$$\begin{aligned} I_{21}(w, w') &= C_h \sum_{k \geq 0} \int_{0 =: u_k < u_{k-1} < u_{k-2} < \dots < u_1 < u_0 \leq t_0} \\ &\quad \times |e^{-X(w')} - e^{-X(w)}| \left(\prod_{i=0}^{k-1} (w'(1, u_i) + C_W) du_i \right) du_{k-1}, \end{aligned}$$

with

$$X(w) = \sum_{i=0}^{k-1} \Omega_{\theta, w, z}(u_{i+1}, u_i) + \Omega_{\theta, w, z}(u_0, t),$$

and

$$\begin{aligned} I_{22}(w, w') &= C_h e^{-\tilde{\Omega}_w(0, t) + C_W t} \sum_{k \geq 1} \int_{0 =: u_k < u_{k-1} < u_{k-2} < \dots < u_1 < u_0 \leq t_0} \\ &\quad \times \left| \prod_{i=0}^{k-1} \tilde{w}_{\theta, w', z}(u_{i+1}, u_i) - \prod_{i=0}^{k-1} \tilde{w}_{\theta, w, z}(u_{i+1}, u_i) \right| \prod_{i=0}^{k-1} du_i. \end{aligned}$$

Note that (78) with (25) implies

$$|\Omega_{\theta,w',z}(u, v) - \Omega_{\theta,w,z}(u, v)| \leq \|w' - w\|_T (v - u), \quad 0 \leq u \leq T.$$

Using this, (83), and (81) in $I_{21}(w, w')$, we further have

$$\begin{aligned} I_{21}(w, w') &\leq C_h e^{-\tilde{\Omega}_w(0,t)+C_W t} e^{\|w' - w\|_T t} \|w' - w\|_T t e^{\tilde{\Omega}_{w'}(0,t)+C_W t} \\ &\leq C_h e^{2C_W T} e^{\|w' - w\|_T T} \|w' - w\|_T T. \end{aligned}$$

With a similar argument, we also have

$$\begin{aligned} I_{22}(w, w') &\leq C_h e^{-\tilde{\Omega}_w(0,t)+C_W t} \sum_{k \geq 1} \sum_{j=0}^{k-1} \int_{0=:u_k < u_{k-1} < u_{k-2} < \dots < u_1 < u_0 \leq t_0} \\ &\quad \times \left(\prod_{i=0}^{j-1} \tilde{w}_{\theta,w',z}(u_{i+1}, u_i) \right) |\tilde{w}_{\theta,w',z}(u_{j+1}, u_j) - \tilde{w}_{\theta,w,z}(u_{j+1}, u_j)| \\ &\quad \times \left(\prod_{i=j+1}^{k-1} \tilde{w}_{\theta,w,z}(u_{i+1}, u_i) \right) \prod_{i=0}^{k-1} du_i \\ &\leq C_h e^{-\tilde{\Omega}_w(0,t)+C_W t} \|w' - w\|_T \int_0^{t_0} e^{\tilde{\Omega}_w(0,v)+C_W v} e^{\tilde{\Omega}_{w'}(v,t_0)+C_W(t_0-v)} dv \\ &\leq C_h e^{-\tilde{\Omega}_w(0,t)+C_W t} \|w' - w\|_T \\ &\quad \times \left(e^{\tilde{\Omega}_w(0,t_0)} t_0 + \int_0^{t_0} e^{\tilde{\Omega}_{w'}(0,v)} |e^{\tilde{\Omega}_{w'}(v,t_0)} - e^{\tilde{\Omega}_w(v,t_0)}| dv \right) \\ &\leq C_h e^{2C_W T} \|w' - w\|_T T (1 + e^{\|w' - w\|_T T} \|w' - w\|_T T). \end{aligned}$$

Since the right hand sides of the bounds for I_{21} and I_{22} are uniform in t_0 and t , these prove equicontinuity of H in $w \in W$.

This completes a proof of equicontinuity of H , hence a proof of Theorem 3.2. \square

4 Hierarchy of multi time Gronwall inequality.

The following is a simple form of Gronwall's inequality.

PROPOSITION 4.1. *Let T be a positive constant, and a and c be non-negative constants. If $x : [0, T] \rightarrow \mathbb{R}$ is an integrable function, satisfying*

$$x(t) \leq a + c \int_0^t x(s) ds, \quad t \in [0, T],$$

then

$$(84) \quad x(t) \leq a e^{ct}, \quad t \in [0, T],$$

holds.

The following is a generalization of Proposition 4.1 to functions of more than 1 variables, where the case $q = 1$ is Proposition 4.1.

PROPOSITION 4.2. *Let T be a positive constant, q a positive integer, and a and c non-negative constants. If $x : [0, T]^q \rightarrow \mathbb{R}$ is an integrable function of q variables, satisfying*

$$x(t_1, \dots, t_q) \leq a e^{c(t_1 + \dots + t_q)} \frac{1}{q} \sum_{i=1}^q e^{-ct_i} + \frac{c}{q} \sum_{i=1}^q \int_0^{t_i} (x(t_1, \dots, t_q)|_{t_i=u}) du, \quad (t_1, \dots, t_q) \in [0, T]^q,$$

then

$$(85) \quad x(t_1, \dots, t_q) \leq a e^{c(t_1 + \dots + t_q)}, \quad (t_1, \dots, t_q) \in [0, T]^q,$$

holds.

To prove Proposition 4.2, we start with the homogeneous case.

PROPOSITION 4.3. *Let T be a positive constant, q a positive integer, and c a non-negative constant. If $x : [0, T]^q \rightarrow \mathbb{R}$ is an integrable function of q variables, satisfying*

$$(86) \quad x(t_1, \dots, t_q) \leq c \sum_{i=1}^q \int_0^{t_i} (x(t_1, \dots, t_q)|_{t_i=s}) ds, \quad (t_1, \dots, t_q) \in [0, T]^q,$$

then

$$(87) \quad x(t_1, \dots, t_q) \leq 0, \quad (t_1, \dots, t_q) \in [0, T]^q,$$

holds.

To prove Proposition 4.3, we introduce a notation

$$(88) \quad \begin{aligned} & (A_{i,k}y)(t_1, \dots, t_q) \\ &= \begin{cases} \frac{1}{(k-1)!} \int_0^{t_i} (t_i - s)^{k-1} (y(t_1, \dots, t_q))|_{t_i=s} ds, & k = 1, 2, 3, \dots, \\ y(t_1, \dots, t_q) \quad (\text{i.e., } A_{i,0} = \text{id}), & k = 0, \end{cases} \end{aligned}$$

for integrable function $y : [0, T]^q \rightarrow \mathbb{R}$ in q variables and $i = 1, \dots, q$. $A_{i,k}$, $k \in \mathbb{Z}_+$, $i = 1, 2, \dots, q$, are commutative operators on the set of integrable functions. In fact, commutativity is obvious for $k = 0$, and by induction in k we have

$$(89) \quad A_{i,k}A_{i,\ell} = A_{i,1}^{k+\ell} = A_{i,k+\ell} = A_{i,\ell}A_{i,k},$$

and Fubini's theorem implies for $k\ell > 0$ and $i \neq j$

$$\begin{aligned} & (A_{i,k}A_{j,\ell}y)(t_1, \dots, t_q) \\ &= \frac{1}{(k-1)!} \frac{1}{(\ell-1)!} \int_0^{t_i} ds \int_0^{t_j} du (t_i - s)^{k-1} \\ & \quad \times ((t_j - u)^{\ell-1} (y(t_1, \dots, t_q))|_{t_j=u})|_{t_i=s} \\ &= (A_{j,\ell}A_{i,k}y)(t_1, \dots, t_q), \end{aligned}$$

which prove

$$(90) \quad A_{i,k}A_{j,\ell} = A_{j,\ell}A_{i,k}, \quad k, \ell \in \mathbb{Z}_+, \quad i, j \in \{1, \dots, q\}.$$

LEMMA 4.4. *Under the assumptions of Proposition 4.3,*

$$(91) \quad x(t_1, \dots, t_q) \leq c^N \sum_{\substack{(k_1, \dots, k_q) \in \mathbb{Z}_+^q; \\ k_1 + \dots + k_q = N}} (A_{q,k_q} A_{q-1,k_{q-1}} \cdots A_{1,k_1} x)(t_1, \dots, t_q), \quad N \in \mathbb{Z}_+,$$

holds.

PROOF. The case $N = 1$ of (91) is the assumption (86) itself. Assume that (91) holds for some N . Substituting (86) in (91), and noting that sums, integrations, and multiplication of non-negative reals have monotonicity, we have

$$x(t_1, \dots, t_q) \leq c^{N+1} \sum_{i=1}^q \sum_{\substack{(k_1, \dots, k_q) \in \mathbb{Z}_+^q; \\ k_1 + \dots + k_q = N}} (A_{q,k_q} A_{q-1,k_{q-1}} \cdots A_{1,k_1} A_{i,1} x)(t_1, \dots, t_q).$$

Using (89) in the form $A_{i,k_i} A_{i,1} = A_{i,k_i+1}$, we have (91) for N replaced by $N + 1$. \square

Proof of Proposition 4.3. For notational simplicity, put $\vec{t} = (t_1, \dots, t_q)$ in this proof. The operator $A_{i,k}$ in (88) satisfies

$$(A_{i,k}y)(\vec{t}) \leq \frac{t_i^k}{k!} \sup_{\vec{t} \in [0, T]^q} y(\vec{t}), \quad \vec{t} \in [0, T]^q,$$

for a integrable function y , hence (91) implies

$$(92) \quad x(\vec{t}) \leq c^N \sum_{\substack{(k_1, \dots, k_q) \in \mathbb{Z}_+^q; \\ k_1 + \dots + k_q = N}} \prod_{i=1}^q \frac{t_i^{k_i}}{k_i!} \sup_{\vec{t} \in [0, T]^q} x(\vec{t}), \quad \vec{t} \in [0, T]^q, \quad N \in \mathbb{N}.$$

For an arbitrary $\epsilon > 0$, let $\vec{t}_0 = (t_{0,1}, \dots, t_{0,q}) \in [0, T]^q$ be a vector (independent of N) such that $x(\vec{t}_0) \geq \sup_{\vec{t} \in [0, T]^q} x(\vec{t}) - \epsilon$ holds. Put

$$a_N = c^N \sum_{\substack{(k_1, \dots, k_q) \in \mathbb{Z}_+^q; \\ k_1 + \dots + k_q = N}} \prod_{i=1}^q \frac{t_{0,i}^{k_i}}{k_i!}.$$

Then (92) implies

$$\sup_{\vec{t} \in [0, T]^q} x(\vec{t}) \leq x(\vec{t}_0) + \epsilon \leq a_N \sup_{\vec{t} \in [0, T]^q} x(\vec{t}) + \epsilon,$$

hence

$$\sup_{\vec{t} \in [0, T]^q} x(\vec{t}) (1 - a_N) \leq \epsilon$$

holds. We see

$$\sum_{N=0}^{\infty} a_N \leq \sum_{N=0}^{\infty} \sum_{\substack{(k_1, \dots, k_q) \in \mathbb{Z}_+^q \\ k_1 + \dots + k_q = N}} \prod_{i=1}^q \frac{(ct_{0,i})^{k_i}}{k_i!} = \prod_{i=1}^q e^{ct_{0,i}} < \infty,$$

so that, in particular, $\lim_{N \rightarrow \infty} a_N = 0$, which implies $1 - a_N \geq \frac{1}{2}$ for large N . Hence

$$\sup_{\vec{t} \in [0, T]^q} x(\vec{t}) \leq 2\epsilon,$$

which proves (87). □

Proof of Proposition 4.2. Note that

$$(93) \quad x_1(s_1, \dots, s_q) = a e^{c(s_1 + \dots + s_q)}$$

satisfies

$$(94) \quad \begin{aligned} x_1(s_1, \dots, s_q) &= e^{-cs_i} x_1(s_1, \dots, s_q) + c \int_0^{s_i} (x_1(s_1, \dots, s_q)|_{s_i=u}) du, \\ (s_1, \dots, s_q) &\in [0, t]^q, \quad i = 1, 2, \dots, q. \end{aligned}$$

Subtracting $x_1(s_1, \dots, s_q)$ from (85), and then using (94), we have

$$\begin{aligned} x(t_1, \dots, t_q) - x_1(t_1, \dots, t_q) &\leq \frac{c}{q} \sum_{i=1}^q \int_0^{t_i} (x(t_1, \dots, t_q) - x_1(t_1, \dots, t_q))|_{t_i=s} ds, \\ (t_1, \dots, t_q) &\in [0, T]^q, \end{aligned}$$

which, with Proposition 4.3 and (93) implies (85). □

Finally, we give a result to be used in the proof of the main theorem in § 5 which contains recursion with respect to the number of variables q and a nonlinear term.

THEOREM 4.5. *Let T be a positive constant, d be a non-negative constant satisfying $d \leq 1$, and for each positive integer q let a_q , b_q , and c_q be non-negative constants. Assume that, for a series of non-negative valued integrable functions $x_q : [0, T]^q \rightarrow [0, \infty)$, $q \in \mathbb{Z}_+$,*

$$(95) \quad \begin{aligned} x_0 &= 1, \\ x_q(t_1, \dots, t_q) &\leq a_q \sum_{i=1}^q x_{q-1}(t_1, \dots, t'_i, \dots, t_q)^d \\ &\quad + b_q \sum_{i=1}^q x_{q-1}(t_1, \dots, t'_i, \dots, t_q) \\ &\quad + c_q \sum_{i=1}^q \int_0^{t_i} (x_q(t_1, \dots, t_q)|_{t_i=s}) ds, \\ (t_1, \dots, t_q) &\in [0, T]^q, \quad q \in \mathbb{N}, \end{aligned}$$

hold. Here, for a function x_{q-1} in $q-1$ variables and $q-1$ variables

$$t_1, t_2, \dots, t_{i-1}, t_{i+1}, \dots, t_q,$$

we wrote

$$x_{q-1}(t_1, \dots, t'_i, \dots, t_q) = x_{q-1}(t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_q).$$

Put

$$\tilde{c}_q = \max_{1 \leq k \leq q} kc_k, \quad q \in \mathbb{N},$$

and define a sequence of non-negative constants g_q , $q = 0, 1, 2, \dots$, recursively by

$$g_0 = 1, \quad g_q = q(a_q g_{q-1}^d + b_q g_{q-1}), \quad q \in \mathbb{N}.$$

Then

$$(96) \quad x_q(t_1, \dots, t_q) \leq g_q e^{\tilde{c}_q(t_1 + \dots + t_q)}, \quad (t_1, \dots, t_q) \in [0, T]^q, \quad q \in \mathbb{N},$$

holds.

PROOF. If $q = 1$, (95) reads $x_1(t_1) \leq (a_1 + b_1) + c_1 \int_0^{t_1} x_1(s) ds$, hence Proposition 4.1 implies $x_1(t_1) \leq g_1 e^{c_1 t_1}$, which proves (96) for $q = 1$.

Let $q \geq 2$ and assume that (96) holds for x_{q-1} , as

$$x_{q-1}(t_1, \dots, t_{q-1}) \leq g_{q-1} e^{\tilde{c}_{q-1}(t_1 + \dots + t_{q-1})}.$$

This and (95) for x_q and $d \leq 1$ imply

$$\begin{aligned} x_q(s_1, \dots, s_q) &\leq q(a_q g_{q-1}^d + b_q g_{q-1}) e^{\tilde{c}_{q-1}(s_1 + \dots + s_q)} \frac{1}{q} \sum_{i=1}^q e^{-\tilde{c}_{q-1} s_i} \\ &\quad + qc_q \frac{1}{q} \sum_{i=1}^q \int_0^{s_i} (x_q(s_1, \dots, s_q)|_{s_i=u}) du, \end{aligned}$$

which, with Proposition 4.2, implies (96) for x_q . □

5 Proof of the main theorem.

5.1 Convergence of the spatial distribution function.

Here we will prove the essential part of the infinite particle limit, the convergence of spatial distribution function.

In analogy to (36), define, for each $i = 1, 2, \dots, N$ and $0 \leq t_0 \leq t \leq T$ and $0 \leq y_0 \leq 1$,

$$(97) \quad J_i^{(N)}(t_0, t) = \{\omega \in \Omega \mid \tilde{v}_i^{(N)}(t)(\omega) > \tilde{v}_i^{(N)}(t_0)(\omega)\}.$$

By similar arguments as for (48) and (38), $\varphi^{(N)}$ in (6) and $Y_C^{(N)}$ in (7) respectively satisfies

$$(98) \quad \varphi^{(N)}(dw, (y_0, t_0), t) = \frac{1}{N} \sum_{j; Y_j^{(N)}(t_0) \geq y_0} \mathbf{1}_{J_j^{(N)}(t_0, t)^c} \delta_{w_j}(dw),$$

and

$$(99) \quad Y_C^{(N)}((y_0, t_0), t) = y_0 + \frac{1}{N} \sum_{j; Y_j^{(N)}(t_0) \geq y_0} \mathbf{1}_{J_j^{(N)}(t_0, t)}.$$

PROPOSITION 5.1. *Assume (8), (20), (24), and (27). Then there exists $\delta' > 0$ and an integer p_0 satisfying $2p_0\delta' > 1$, such that for any integer $p \geq p_0$ there exists a positive constant C depending only on p and δ' , (independent of N and h), such that for any bounded continuous $h : W \rightarrow \mathbb{R}$*

$$(100) \quad \mathbb{E} \left[\sup_{(\gamma, t) \in \Delta_T} \left| \varphi^{(N)}(h, \gamma, t) - \varphi_{y_C}(h, \gamma, t) \right|^{2p} \right] \leq \frac{C C_h^{2p}}{N^{2p\delta'}}, \quad N \in \mathbb{N},$$

holds, where C_h is as in (51).

Note that (7) and (17) with $\theta = y_C$ imply

$$Y_C^{(N)}(\gamma, t) - y_C(\gamma, t) = \frac{[N(1 - y_0)]}{N} - (1 - y_0) + \varphi_{y_C}(W, \gamma, t) - \varphi^{(N)}(W, \gamma, t).$$

Applying $(a + b)^{2p} \leq 2^{2p-1}(a^{2p} + b^{2p})$, valid for $a, b \geq 0$ and $2p \geq 1$, Proposition 5.1 with $h(w) = 1$, $w \in W$, therefore implies

$$(101) \quad \mathbb{E} \left[\sup_{(\gamma, t) \in \Delta_T} \left| Y_C^{(N)}(\gamma, t) - y_C(\gamma, t) \right|^{2p} \right] \leq \frac{2^{2p-1}C}{N^{2p\delta'}} + \frac{2^{2p-1}}{N^{2p}}, \quad N \in \mathbb{N},$$

with the assumptions and notations of the Theorem.

5.1.1 Coupling of the original and the flow driven model.

In view of Theorem 3.2, it suffices to prove the following for Proposition 5.1 to hold.

PROPOSITION 5.2. *Assume (8), (20), (24), and (27), and let δ be as in (20). Then there exists $\delta' > 0$ and an integer p_0 satisfying $2p_0\delta' > 1$, such that for any integer $p \geq p_0$ there exists a positive constant C , (independent of N and h ,) such that for any bounded continuous $h : W \rightarrow \mathbb{R}$*

$$(102) \quad \mathbb{E} \left[\sup_{((y_0, t_0), t) \in \Delta_T} \left| \varphi^{(N)}(h, \gamma, t) - \varphi^{(N, y_C)}(h, \gamma, t) \right|^{2p} \right] \leq \frac{C C_h^{2p}}{N^{2p\delta'}}, \quad N \in \mathbb{N},$$

where C_h is as in (51).

For $t \in [0, T]$ and $i \in \{1, 2, \dots, N\}$, put

$$(103) \quad \tilde{w}_{i, \wedge}^{(N)}(t) = w_i(Y_i^{(N)}(t-), t) \wedge w_i(y_C(\gamma_i^{(N, y_C)}(t-), t), t),$$

and

$$(104) \quad \tilde{w}_{i, \vee}^{(N)}(t) = w_i(Y_i^{(N)}(t-), t) \vee w_i(y_C(\gamma_i^{(N, y_C)}(t-), t), t),$$

and denote the event that the i -th particle $Y_i^{(N)}(s)$ of (1) and $Y_i^{(N, y_C)}(s)$ of (35) jump to top at same times in the interval $(t_0, t]$ by

$$(105) \quad \mathcal{K}_i^{(N)}(t_0, t) = \{\omega \in \Omega \mid \nu_i^{(N)}(\{(s, \xi) \mid \tilde{w}_{i, \wedge}^{(N)}(s) < \xi \leq \tilde{w}_{i, \vee}^{(N)}(s), s \in (t_0, t]\}) = 0\}.$$

The definitions (98) and (53) of $\varphi^{(N)}$ and $\varphi^{(N, \theta)}$ are defined for $(y_0, t_0) \in [0, 1] \times [0, t]$. Note that if we restrict (y_0, t_0) to the initial/boundary points Γ_t of (10), then the summation over j in these definitions are equivalent to the summation over j satisfying $y_j^{(N)} \geq y_0$. In particular, the summations are deterministic and same for both $\varphi^{(N)}$ and $\varphi^{(N, \theta)}$, if $(y_0, t_0) \in \Gamma_t$. This is because $(y_0, t_0) \in \Gamma_t$ implies $y_0 = 0$ or $t_0 = 0$, by (10), and if $y_0 = 0$ then the condition $Y_j^{(N)}(t_0) \geq y_0$ (or $Y_j^{(N, \theta)}(t_0) \geq y_0$) in (98) ((53), respectively), implies summation over all j , while if $t_0 = 0$ then (1) and (31) imply $Y_j^{(N)}(t_0) = Y_j^{(N, \theta)}(t_0) = y_j^{(N)}$. This proves that for all the cases of $\varphi^{(N)}$ and $\varphi^{(N, \theta)}$ with $(y_0, t_0) \in \Gamma_t$ the summation over j are equivalent to the the summation over j satisfying $y_j^{(N)} \geq y_0$.

Fix a bounded continuous function $h : W \rightarrow \mathbb{R}$, and let C_h be as in (51). Using the definitions (98) and (53), put

$$(106) \quad \begin{aligned} \Delta\varphi^{(N)}(\gamma, t) &= \varphi^{(N)}(h, \gamma, t) - \varphi^{(N, \theta)}(h, \gamma, t) \\ &= \frac{1}{N} \sum_{j; y_j^{(N)} \geq y_0} h(w_j)(\mathbf{1}_{J_j^{(N)}(t_0, t)^c} - \mathbf{1}_{J_j^{(N, \theta)}(t_0, t)^c}), \\ \gamma &= (y_0, t_0) \in \Gamma_t, \quad t \in [0, T]. \end{aligned}$$

Then (105) and (51) imply,

$$|\Delta\varphi^{(N)}(\gamma, t)| \leq \frac{C_h}{N} \sum_{j=1}^N \mathbf{1}_{\mathcal{K}_j^{(N)}(t_0, t)^c}, \quad \gamma = (y_0, t_0) \in \Gamma_t, \quad t \in [0, T].$$

The monotonicity of $\mathcal{K}_j^{(N)}(t_0, t)^c$ with respect to t and t_0 further implies

$$(107) \quad \sup_{(\gamma, t) \in \Delta_T} |\Delta\varphi^{(N)}(\gamma, t)| \leq \frac{C_h}{N} \sum_{i=1}^N \mathbf{1}_{\mathcal{K}_i^{(N)}(0, T)^c},$$

Proof of Proposition 5.2 therefore reduces to evaluation of the event $\mathcal{K}_i^{(N)}(0, T)$.

5.1.2 Event with different jumps to top.

As an analog of (32), define a sequence of stopping times, $0 = \tau_{i,0}^{(N)} < \tau_{i,1}^{(N)} < \dots$, by

$$(108) \quad \begin{aligned} \tau_{i,0}^{(N)} &= 0, \\ \tau_{i,k+1}^{(N)} &= \inf\{t > \tau_{i,k}^{(N)} \mid \nu_i^{(N)}(\{(s, \xi) \in (\tau_{i,k}^{(N)}, T] \times [0, \infty) \mid \\ &\quad 0 \leq \xi \leq w_i(Y_i^{(N)}(s-), s)\}) > 0\}, \quad k \in \mathbb{Z}_+. \end{aligned}$$

$\tau_{i,k}^{(N)}$ is the time that the particle i in the (original) stochastic ranking process jumps to the top for the k -th time. A corresponding analog of (33) is

$$(109) \quad \gamma_i^{(N)}(t) = \begin{cases} (y_i^{(N)}, 0), & 0 \leq t < \tau_{i,1}^{(N)}, \\ (0, \tau_{i,k}^{(N)}), & \tau_{i,k}^{(N)} \leq t < \tau_{i,k+1}^{(N)}, \quad k = 1, 2, \dots \end{cases}$$

A property corresponding to (42) then is

$$(110) \quad Y_i^{(N)}(t) = Y_C^{(N)}(\gamma_i^{(N)}(t), t), \quad t \in [0, T],$$

which can be proved in a similar way as a proof of (42) in Lemma 3.1. This decomposition in particular decomposes the dependence as random variables; if we temporarily denote by $X \in \mathcal{F}$, a fact that a random variable $X : \Omega \rightarrow \mathbb{R}$ is \mathcal{F} -measurable, and denote by $\sigma[Z]$ a sigma algebra generated by a random variable Z , we have

$$(111) \quad \begin{aligned} Y_C^{(N)}((y_0, t_0), t) &\in \sigma[\{\nu_j^{(N)} \mid Y_j^{(N)}(t_0) > y_0\}], \\ \gamma_i^{(N)} &\in \sigma[\{\tau_{i,k}^{(N)} \wedge t \mid k \in \mathbb{N}\}]. \end{aligned}$$

Define an analog of the stopping times (108) using (103) by

$$(112) \quad \begin{aligned} \tau_{i,\wedge,0}^{(N)} &= 0, \\ \tau_{i,\wedge,k}^{(N)} &= \inf\{t > \tau_{i,\wedge,k-1}^{(N)} \mid \nu_i^{(N)}(\{(\xi, s) \mid 0 \leq \xi \leq \tilde{w}_{i,\wedge}^{(N)}(s), \quad 0 \leq s \leq t\}) > 0\}, \\ &k \in \mathbb{N}, \end{aligned}$$

and denote by $\sigma_i^{(N)}$, the time that the particle pair with label i of the original model and the flow driven model have different jumps to the top for the first time;

$$(113) \quad \sigma_i^{(N)}(\omega) = \inf\{t \in [0, T] \mid \omega \in \mathcal{K}_i^{(N)}(0, t)^c\}, \quad \omega \in \Omega.$$

The definition implies

$$(114) \quad \tau_{i,k}^{(N)} < \sigma_i^{(N)} \Rightarrow \tau_{i,\wedge,k}^{(N)} = \tau_{i,k}^{(N)} = \tau_{i,k}^{(N,y_C)},$$

where $\tau_{i,k}^{(N,y_C)}$ is defined in (32), with $\theta = y_C$. Independence of $\nu_i^{(N)}(A)$ and $\nu_i^{(N)}(B)$ for the exclusive events A and B implies

$$(115) \quad \{\tau_{i,\wedge,k}^{(N)} \mid k \in \mathbb{Z}_+\} \perp \sigma_i^{(N)}.$$

Using (104), (103), (8), and $a \vee b = |a - b| + a \wedge b$, we have

$$(116) \quad \begin{aligned} & \{(\xi, s) \in \mathbb{R}^2 \mid \tilde{w}_{i,\wedge}^{(N)}(s) < \xi \leq \tilde{w}_{i,\vee}^{(N)}(s), 0 < s \leq t\} \\ & \subset \{(\xi, s) \in \mathbb{R}^2 \mid 0 \leq \xi - \tilde{w}_{i,\wedge}^{(N)}(s) \leq C_W |Y_i^{(N)}(s) - y_C(\gamma_i^{(N,y_C)}(s), s)|, \\ & \quad 0 < s \leq t\} \\ & \subset \bigcup_{k=1}^{\infty} \{(\xi, s) \in \mathbb{R}^2 \mid 0 \leq \xi - \tilde{w}_{i,\wedge}^{(N)}(s) \leq C_W |Y_i^{(N)}(s) - y_C(\gamma_i^{(N,y_C)}(s), s)|, \\ & \quad t \wedge \tau_{i,\wedge,k-1}^{(N)} < s \leq t \wedge \tau_{i,\wedge,k}^{(N)}\}. \end{aligned}$$

Note that for each i

$$Y_i^{(N)}(\tau_{i,k-1}^{(N)}) = Y_i^{(N,y_C)}(\tau_{i,k-1}^{(N)}) = 0, \quad \text{on } \tau_{i,k-1}^{(N)} < \sigma_i^{(N)}.$$

Note also that the definition (113) implies $\{\omega \in \Omega \mid \sigma_i^{(N)}(\omega) > s\} = \mathcal{K}_i^{(N)}(0, s)$. Hence, (110), (42), (99), and (38) imply, with similar arguments for deriving (107) from (105),

$$\begin{aligned} |Y_i^{(N)}(s) - Y_i^{(N,y_C)}(s)| &= |Y_C^{(N)}(\gamma_i^{(N)}(s), s) - Y_C^{(N,y_C)}(\gamma_i^{(N)}(s), s)| \\ &= \frac{1}{N} \left| \sum_{j \neq i} (\mathbf{1}_{(J_j^{(N)}(\tau_{i,k-1}^{(N)}, s))^c} - \mathbf{1}_{(J_j^{(N)}(\tau_{i,k-1}^{(N,y_C)}, s))^c}) \right| \\ &\leq \frac{1}{N} \sum_{j \neq i} \mathbf{1}_{\mathcal{K}_j^{(N)}(\tau_{i,k-1}^{(N)}, s)^c} \\ &\text{on } \mathcal{K}_i^{(N)}(0, s), \quad \tau_{i,k-1}^{(N)} \leq s < \tau_{i,k}^{(N)}. \end{aligned}$$

Hence up to the first jump to the top, each $Y_{i_\alpha}^{(N)}(t)$ with $i_\alpha \neq i_0$ is independent of ν_{i_0} . Therefore,

$$\begin{aligned} & \mathbb{E} \left[\prod_{\alpha=1}^q \mathbf{1}_{\mathcal{K}_{i_\alpha}^{(N)}(0, t_\alpha)^c} \right] \\ &= \mathbb{E} \left[\prod_{\alpha; i_\alpha \neq i_0} \mathbf{1}_{(\mathcal{K}_{i_\alpha}^{(N)}(0, t_\alpha))^c} \mathbb{P} \left[(\mathcal{K}_{i_0}^{(N)}(0, t_0))^c \mid \{\nu_j, j \neq i_0\} \cup \{\tau_{i_0, \wedge, k}^{(N)}\} \right] \right], \end{aligned}$$

where $\mathbb{P}[\cdot \mid \{\nu_j, j \neq i_0\} \cup \{\tau_{i_0, \wedge, k}^{(N)}\}]$ denotes conditional probability conditioned on the sigma algebra generated by $\nu_j, j \neq i_0$, and $\tau_{i_0, \wedge, k}^{(N)}, k \in \mathbb{N}$. This with (115), (114), (117), and

$$\mathbb{P}[\nu(A) > 0] = 1 - e^{-|A|} \leq |A|$$

for a unit Poisson random measure ν , further leads to

$$\begin{aligned} & \mathbb{E} \left[\prod_{\alpha=1}^q \mathbf{1}_{\mathcal{K}_{i_\alpha}^{(N)}(0, t_\alpha)^c} \right] \\ & \leq \mathbb{E} \left[\prod_{\alpha; i_\alpha \neq i_0} \mathbf{1}_{(\mathcal{K}_{i_\alpha}^{(N)}(0, t_\alpha))^c} \sum_{k=1}^{\infty} \mathbf{1}_{\mathcal{K}_{i_0}^{(N)}(0, t_0 \wedge \tau_{i_0, \wedge, k-1}^{(N)})} \right. \\ & \quad \times \left(\int_{t_0 \wedge \tau_{i_0, \wedge, k-1}^{(N)}}^{t_0 \wedge \tau_{i_0, \wedge, k}^{(N)}} C_W |Y_{i_0}^{(N, y_C)}(s) - y_C(\gamma_{i_0}^{(N, y_C)}(s), s)| ds \right. \\ & \quad \left. \left. + \int_{t_0 \wedge \tau_{i_0, \wedge, k-1}^{(N)}}^{t_0 \wedge \tau_{i_0, \wedge, k}^{(N)}} \frac{C_W}{N} \sum_{j \neq i_0} \mathbf{1}_{\mathcal{K}_j^{(N)}(t_0 \wedge \tau_{i_0, \wedge, k-1}^{(N)}, s)^c} ds \right) \right] \\ (121) \quad & \leq \frac{C_W}{N} \sum_{j=1}^N \mathbb{E} \left[\prod_{\alpha; i_\alpha \neq i_0} \mathbf{1}_{(\mathcal{K}_{i_\alpha}^{(N)}(0, t_\alpha))^c} \sum_{k=1}^{\infty} \mathbf{1}_{\mathcal{K}_{i_0}^{(N)}(0, t_0 \wedge \tau_{i_0, \wedge, k-1}^{(N)})} \right. \\ & \quad \times \left. \int_{t_0 \wedge \tau_{i_0, \wedge, k-1}^{(N)}}^{t_0 \wedge \tau_{i_0, \wedge, k}^{(N)}} \mathbf{1}_{\mathcal{K}_j^{(N)}(t_0 \wedge \tau_{i_0, \wedge, k-1}^{(N)}, s-)^c} ds \right] \\ & \quad + C_W \mathbb{E} \left[\prod_{\alpha; i_\alpha \neq i_0} \mathbf{1}_{(\mathcal{K}_{i_\alpha}^{(N)}(0, t_\alpha))^c} \sum_{k=1}^{\infty} \mathbf{1}_{\mathcal{K}_{i_0}^{(N)}(0, t_0 \wedge \tau_{i_0, \wedge, k-1}^{(N)})} \right. \\ & \quad \times \left. \int_{t_0 \wedge \tau_{i_0, \wedge, k-1}^{(N)}}^{t_0 \wedge \tau_{i_0, \wedge, k}^{(N)}} |Y_{i_0}^{(N, y_C)}(s) - y_C(\gamma_{i_0}^{(N, y_C)}(s), s)| ds \right]. \end{aligned}$$

Using $\mathbf{1}_{\mathcal{K}_{i_0}^{(N)}(0, t_0 \wedge \tau_{i_0, \wedge, k-1}^{(N)})} \leq 1$ and $\mathbf{1}_{\mathcal{K}_j^{(N)}(t_0 \wedge \tau_{i_0, \wedge, k-1}^{(N)}, s-)^c} \leq \mathbf{1}_{\mathcal{K}_j^{(N)}(0, s-)^c}$, we see that for $i, j = 1, 2, \dots, N$,

$$(122) \quad \sum_{k \geq 1} \mathbf{1}_{\mathcal{K}_i^{(N)}(0, t \wedge \tau_{i, k-1}^{(N)})} \int_{t \wedge \tau_{i, k-1}^{(N)}}^{t \wedge \tau_{i, k}^{(N)}} \mathbf{1}_{\mathcal{K}_j^{(N)}(t \wedge \tau_{i, k-1}^{(N)}, s-)^c} ds \leq \int_0^t \mathbf{1}_{\mathcal{K}_j^{(N)}(0, s-)^c} ds.$$

Substituting (122) in the first term of the right hand side of (121), and bounding the characteristic function on the right hand side by 1 for $j \in \{i_1, \dots, i_q\} \setminus \{i_0\}$, and bounding

the characteristic function for i_0 also by 1 in the second term of the right hand side of (121), we have

$$\begin{aligned}
& X_q^{(N)}(t_1, \dots, t_q) \\
& \leq \frac{C_W T(q-1)}{N} \sum_{i_0=1}^q X_{q-1}^{(N)}(t_1, \dots, t_{i_0}, \dots, t_q) \\
(123) \quad & + C_W \sum_{i_0=1}^q \int_0^{t_{i_0}} X_q^{(N)}(t_1, \dots, s, \dots, t_q) ds \\
& + C_W \max_{\{i_1, \dots, i_q\} \subset \{1, \dots, N\}} \int_0^{t_{i_0}} \\
& \mathbb{E} \left[\prod_{i_\alpha \neq i_0} \mathbf{1}_{\mathcal{K}_{i_\alpha}(0, t_\alpha)^c} |Y_{i_{i_0}}^{(N, y_C)}(s) - y_C((\gamma_{i_{i_0}}^{(N, y_C)}(s), s))| \right] ds.
\end{aligned}$$

Here, $X_{q-1}^{(N)}(t_1, \dots, t_{i_0}, \dots, t_q)$ is the function in (118) with q replaced by $q-1$ and with $q-1$ variables obtained by excluding t_{i_0} from t_1, \dots, t_q , and the variables for $X_q^{(N)}(t_1, \dots, s, \dots, t_q)$ is t_1, \dots, t_q with t_{i_0} replaced by s . Applying Hölder's inequality in the form

$$\mathbb{E}[|XY|] \leq \mathbb{E}[|X|^{2p/(2p-1)}]^{1-(2p)^{-1}} \mathbb{E}[|Y|^{2p}]^{1/(2p)}$$

to the last term in the right hand side of (123), and using (42) and (56), we have

$$\begin{aligned}
& X_q^{(N)}(t_1, \dots, t_q) \\
& \leq \frac{C_W T(q-1)}{N} \sum_{i_0=1}^q X_{q-1}^{(N)}(t_1, \dots, t_{i_0}, \dots, t_q) \\
& + C_W \sum_{i_0=1}^q \int_0^{t_{i_0}} X_q^{(N)}(t_1, \dots, s, \dots, t_q) ds \\
& + \frac{C_W TC}{N^\delta} \sum_{i_0=1}^q (X_{q-1}^{(N)}(t_1, \dots, t_{i_0}, \dots, t_q))^{(2p-1)/(2p)}, \\
& q = 1, 2, \dots, \\
& X_0^{(N)} = 1.
\end{aligned}$$

Applying Theorem 4.5, with $a_q = C_W TC N^{-\delta}$, $b_q = C_W T(q-1)N^{-1}$, $c_q = C_W$, $d = 1 - (2p)^{-1}$, we have

$$\begin{aligned}
(124) \quad & X_q^{(N)}(t_1, \dots, t_q) \leq g_q e^{q^2 C_W T}, \quad t_i \in [0, T], \quad i = 1, \dots, q, \quad q \in \mathbb{N}; \\
& g_0 = 1, \quad g_q = q C_W T (C N^{-\delta} g_{q-1}^d + (q-1)N^{-1} g_{q-1}), \quad q \in \mathbb{N}.
\end{aligned}$$

For large N we have $g_{q-1} < 1$, hence with $d < 1$, we further have $g_{q-1}^d > g_{q-1}$, and also with $0 < \delta < 1$ for large N we have $N^{-\delta} > (q-1)N^{-1}$, so that

$$g_q \leq q C_W T g_{q-1}^d N^{-\delta} (C+1), \quad q \in \mathbb{N}.$$

By induction in q ,

$$g_q \leq q!(C_W T \vee 1)^q (C + 1)^q \frac{1}{N^{2p\delta(1-(1-(2p)^{-1})^q)}}, \quad q \in \mathbb{N}.$$

Since $1 - (1 - \frac{1}{2p})^q$ is decreasing in q , we therefore have

$$(125) \quad \begin{aligned} X_q^{(N)}(t_1, \dots, t_q) &\leq q!(C_W T \vee 1)^q (C + 1)^q e^{q^2 C_W T} \frac{1}{N^{2p\delta(1-(1-(2p)^{-1})^{2p})}}, \\ q &= 1, 2, \dots, p. \end{aligned}$$

Choose δ' to be any positive constant satisfying

$$0 < \delta' < (1 - \frac{1}{e})\delta.$$

Since $\lim_{p \rightarrow \infty} (1 - \frac{1}{2p})^{2p} = e^{-1} < 1$, there exists an integer $p_0 > \frac{1}{2\delta'}$ such that

$$\delta' < (1 - (1 - \frac{1}{2p})^{2p})\delta, \quad p = p_0, p_0 + 1, \dots$$

With (125) we arrive at

$$X_q^{(N)}(t_1, \dots, t_q) \leq q!(C_W T \vee 1)^q (C + 1)^q e^{q^2 C_W T} \frac{1}{N^{2p\delta'}}, \quad q = 1, 2, \dots, 2p,$$

for $p = p_0, p_0 + 1, \dots$. Since $\frac{N!}{q!(N-q)!} \leq \frac{N^q}{q!} \leq \frac{N^{2p}}{q!}$, this proves (119).

This completes a proof of Proposition 5.2, and therefore, of Proposition 5.1. \square

5.2 Proof of Theorem 2.3.

Let $y \in [0, 1]$ and let $h : W \rightarrow \mathbb{R}$ be a bounded continuous function with $C_h > 0$ as in (51). For the flow $y_C \in \Theta_T$ in Theorem 2.1, the definition of Θ_T in (12) implies that for each $t \in [0, T]$, $\Gamma_t \ni \gamma \mapsto y_C(\gamma, t) \in [0, 1]$ is surjective. Therefore there exists $\gamma_t : [0, 1] \rightarrow \Gamma_t$ such that

$$(126) \quad y_C(\gamma_t(y), t) = y, \quad y \in [0, 1].$$

We then have, using (18), (54), (126), (6), (17), (51), (3), and (126) in turn,

$$\begin{aligned}
& \left| \int_W h(w) \mu_t^{(N)}(dw \times [y, 1]) - \int_W h(w) \mu_t(dw \times [y, 1]) \right| \\
&= \left| \int_W h(w) \mu_t^{(N)}(dw \times [y, 1]) - \varphi_{y_C}(h, \gamma_t(y), t) \right| \\
&\leq \sup_{\gamma \in \Gamma_t} |\varphi^{(N)}(h, \gamma, t) - \varphi_{y_C}(h, \gamma, t)| \\
&\quad + \left| \int_W h(w) \mu_t^{(N)}(dw \times [y, 1]) - \int_W h(w) \mu_t^{(N)}(dw \times [Y_C^{(N)}(\gamma_t(y), t), 1]) \right| \\
&\leq \sup_{\gamma \in \Gamma_t} |\varphi^{(N)}(h, \gamma, t) - \varphi_{y_C}(h, \gamma, t)| + C_h |y - Y_C^{(N)}(\gamma_t(y), t)| \\
&\leq \sup_{\gamma \in \Gamma_t} |\varphi^{(N)}(h, \gamma, t) - \varphi_{y_C}(h, \gamma, t)| + C_h |y_C(\gamma_t(y), t) - Y_C^{(N)}(\gamma_t(y), t)| \\
&\leq \sup_{\gamma \in \Gamma_t} |\varphi^{(N)}(h, \gamma, t) - \varphi_{y_C}(h, \gamma, t)| + C_h \sup_{\gamma \in \Gamma_t} |y_C(\gamma, t) - Y_C^{(N)}(\gamma, t)|.
\end{aligned}$$

Proposition 5.1 and (101) then imply

$$\begin{aligned}
(127) \quad & \mathbb{E} \left[\sup_{t \in [0, T]} \left| \int_W h(w) \mu_t^{(N)}(dw \times [y, 1]) - \int_W h(w) \mu_t(dw \times [y, 1]) \right|^{2p} \right] \\
& \leq \frac{(1 + 2^{2p-1} C_h) C}{N^{2p\delta'}} + \frac{2^{2p-1} C_h}{N^{2p}}, \quad N \in \mathbb{N},
\end{aligned}$$

where the constants in the right hand side is as in Proposition 5.1. Since $2p\delta' > 1$, we see that

$$\mathbb{E} \left[\sum_{N=1}^{\infty} \sup_{t \in [0, T]} \left| \int_W h(w) \mu_t^{(N)}(dw \times [y, 1]) - \int_W h(w) \mu_t(dw \times [y, 1]) \right|^{2p} \right] < \infty,$$

hence, in particular, we have, as in the argument below Corollary 3.3 for Theorem 3.2,

$$(128) \quad \lim_{N \rightarrow \infty} \sup_{t \in [0, T]} \left| \int_W h(w) \mu_t^{(N)}(dw \times [y, 1]) - \int_W h(w) \mu_t(dw \times [y, 1]) \right| = 0,$$

with probability 1, which proves (28).

Next we prove uniform almost sure convergence of $Y_i^{(N)}$ to Y_i for $i = 1, 2, \dots, L$. As an analogy to (32) and (33), define

$$\begin{aligned}
(129) \quad & \tau_{i,0} = 0, \\
& \tau_{i,k+1} = \inf \{ t > \tau_{i,k} \mid \nu_i(\{(s, \xi) \in (\tau_{i,k}, T] \times [0, \infty) \mid \\
& \quad 0 \leq \xi \leq w_i(Y_i(s-), s)\}) > 0 \}, \quad k \in \mathbb{Z}_+,
\end{aligned}$$

and

$$(130) \quad \gamma_i(t) = \begin{cases} (y_i, 0), & 0 \leq t < \tau_{i,1}, \\ (0, \tau_{i,k}), & \tau_{i,k} \leq t < \tau_{i,k+1}, \quad k = 1, 2, \dots \end{cases}$$

Comparing (19) and (30), we have, with similar argument for (42),

$$(131) \quad Y_i(t) = y_C(\gamma_i(t), t), \quad t \in [0, T].$$

Quantities corresponding to (103), (104), (105), and (113) are

$$\begin{aligned} \check{w}_{i,\wedge}^{(N)}(t) &= w_i(Y_i^{(N)}(t-), t) \wedge w_i(Y_i(t-), t), \\ \check{w}_{i,\vee}^{(N)}(t) &= w_i(Y_i^{(N)}(t-), t) \vee w_i(Y_i(t-), t), \\ \tilde{\mathcal{K}}_i^{(N)}(0, t) &= \{\omega \in \Omega \mid \nu_i(\{(s, \xi) \mid \check{w}_{i,\wedge}^{(N)}(s) < \xi \leq \check{w}_{i,\vee}^{(N)}(s), s \in (0, t]\}) = 0\}, \\ \tilde{\sigma}_i^{(N)} &= \inf\{t \in [0, T] \mid \tilde{\mathcal{K}}_i^{(N)}(0, t)^c\}. \end{aligned}$$

A proof now proceeds with argument similar to that in § 5.1.2. An argument similar to that for (116) leads to a bound

$$(132) \quad \begin{aligned} &\tilde{\mathcal{K}}_i^{(N)}(0, t)^c \\ &\subset \{\omega \in \Omega \mid \nu_i(\{(s, \xi) \mid 0 \leq \xi - \check{w}_{i,\wedge}^{(N)}(s) \leq C_W |Y_i^{(N)}(s) - Y_i(s)|, \\ &\quad s \in (0, t]\}) > 0\}. \end{aligned}$$

Since ν_i is a Poisson random measure, it holds with probability 1 that $\nu_i(A) < \infty$ for a Borel set $A \subset [0, T] \times [0, \infty)$ of finite area. Hence for almost all $\omega \in \Omega$ there exists $\epsilon = \epsilon(\omega) > 0$ such that

$$(133) \quad \nu_i(\{(s, \xi) \mid 0 \leq \xi - \check{w}_{i,\wedge}^{(N)}(s) \leq \epsilon, s \in (0, t]\}) = 0.$$

On the other hand, applying the argument from (127) to (128), (101) implies

$$(134) \quad \lim_{N \rightarrow \infty} \sup_{(\gamma, t) \in \Delta_T} |Y_C^{(N)}(\gamma, t) - y_C(\gamma, t)| = 0, \quad a.s..$$

Therefore, for almost all $\omega \in \Omega$, there exists an integer $N_0 = N_0(\omega)$ such that for $N \geq N_0$,

$$(135) \quad \sup_{(\gamma, t) \in \Delta_T} |Y_C^{(N)}(\gamma, t) - y_C(\gamma, t)| \leq \frac{\epsilon}{C_W}, \quad N \geq N_0.$$

Combining (133), (135), (110), (131), and (132),

$$(136) \quad \tilde{\mathcal{K}}_i^{(N)}(0, T)^c = \emptyset, \quad N \geq N_0.$$

Next, (110) and (131) imply

$$\begin{aligned} &|Y_i^{(N)}(t) - Y_i(t)| \\ &\leq |Y_C^{(N)}(\gamma_i^{(N)}(t), t) - y_C(\gamma_i^{(N)}(t), t)| + |y_C(\gamma_i^{(N)}(t), t) - y_C(\gamma_i(t), t)| \\ &\leq \sup_{\gamma \in \Gamma_t} |Y_C^{(N)}(\gamma, t) - y_C(\gamma, t)| + |y_C(\gamma_i^{(N)}(t), t) - y_C(\gamma_i(t), t)|. \end{aligned}$$

Comparing (109) and (130) we further have

$$(137) \quad \begin{aligned} & |Y_i^{(N)}(t) - Y_i(t)| \\ & \leq \sup_{\gamma \in \Gamma_t} |Y_C^{(N)}(\gamma, t) - y_C(\gamma, t)| \\ & \quad + |y_C((y_i^{(N)}, 0), t) - y_C((y_i, 0), t)| \mathbf{1}_{t < \tau_{i,1}} + \mathbf{1}_{\tilde{\mathcal{K}}_i^{(N)}(0,t)^c}. \end{aligned}$$

Substituting (136) in (137),

$$(138) \quad \begin{aligned} & |Y_i^{(N)}(t) - Y_i(t)| \\ & \leq \sup_{\gamma \in \Gamma_t} |Y_C^{(N)}(\gamma, t) - y_C(\gamma, t)| + |y_C((y_i^{(N)}, 0), t) - y_C((y_i, 0), t)|, \quad N \geq N_0. \end{aligned}$$

Since $y_C \in \Theta_T$, $\lim_{N \rightarrow \infty} y_i^{(N)} = y_i$ implies

$$(139) \quad \lim_{N \rightarrow \infty} \sup_{t \in [0, T]} |y_C((y_i^{(N)}, 0), t) - y_C((y_i, 0), t)| = 0, \quad a.s..$$

Combining (134), (138), (139) we have

$$\lim_{N \rightarrow \infty} \sup_{t \in [0, T]} |Y_i^{(N)}(t) - Y_i(t)| = 0, \quad a.s..$$

Therefore the almost sure uniform convergence of tagged particle system holds, which completes a proof of Theorem 2.3.

A Point process with last-arrival-time dependent intensity.

We will summarize the definition and basic formulas of the point processes with last-arrival-time dependent intensities. See [11, §3] and [12, §1.2] for a proof. In accordance with [11, 12], we will denote the point process with last-arrival-time dependent intensity by $N = N(t)$, while we wrote $\tilde{\nu}$ in the main body of the present paper to keep the symbol N for the particle number.

$N = N(t)$, $t \geq 0$, is a non-decreasing, right-continuous, non-negative integer valued stochastic process on a measurable space with $N(0) = 0$, satisfying the following. For each non-negative integer k denote the k -th arrival time τ_k by

$$(140) \quad \tau_k = \inf\{t \geq 0 \mid N(t) \geq k\}, \quad k = 1, 2, \dots, \quad \text{and} \quad \tau_0 = 0.$$

The arrival times τ_k are non-decreasing in k , because N is non-decreasing, and since N is also right-continuous, the arrival times are stopping times; if we denote the associated filtration by $\mathcal{F}_t = \sigma[N(s), s \leq t]$, then $\{\tau_k \leq t\} \in \mathcal{F}_t$, $t \geq 0$. We call N a point

process with last-arrival-time dependent intensities. if for a non-negative valued bounded continuous function ω of (s, t) for $0 \leq s \leq t$, and for $k = 1, 2, \dots$

$$(141) \quad \mathbb{P}[t < \tau_k \mid \mathcal{F}_{\tau_{k-1}}] = \exp\left(-\int_{\tau_{k-1}}^t \omega(\tau_{k-1}, u) du\right) \quad \text{on } t \geq \tau_{k-1}$$

holds.

In particular, (141) with $k = 1$ implies

$$(142) \quad \mathbb{P}[N(t) = 0] = \mathbb{P}[\tau_1 > t] = \exp\left(-\int_0^t \omega(0, u) du\right), \quad t \geq 0.$$

If ω is independent of the first variable, then (141) implies that N is the (inhomogeneous) Poisson process with intensity function ω . We are considering a generalization of the Poisson process such that the intensity function depends on the latest arrival time.

A construction (existence proof) of the point process with last-arrival-time dependent intensity goes as follows. Let $\omega : [0, \infty)^2 \rightarrow [0, \infty)$ be a non-negative valued bounded continuous function of (s, t) for $0 \leq s \leq t$, for which we aim to construct a process satisfying (141). Let ν be a Poisson random measure on $[0, \infty)^2$, with unit constant intensity

$$(143) \quad \mathbb{E}[\nu([a, b] \times [c, d])] = (b - a)(d - c) \quad b > a > 0, \quad d > c > 0, \quad k \in \mathbb{N}.$$

Define a sequence of hitting times $\tilde{\tau}_k$, $k \in \mathbb{Z}_+$, inductively by

$$(144) \quad \begin{aligned} \tilde{\tau}_0 &= 0, \quad \text{and} \\ \tilde{\tau}_k &= \inf\{t \geq \tilde{\tau}_{k-1} \mid \nu(\{(\xi, u) \in [0, \infty)^2 \mid \\ &\quad 0 \leq \xi \leq \omega(\tilde{\tau}_{k-1}, u), \tilde{\tau}_{k-1} < u \leq t\}) > 0\}, \\ &k = 1, 2, \dots \end{aligned}$$

Note that the definition is *not* equivalent to the *wrong* formula such as $\tilde{\tau}_k = \inf\{t \geq 0 \mid \nu(\{(\xi, u) \in [0, \infty)^2 \mid 0 \leq \xi \leq \omega(\tilde{\tau}_{k-1}, u), 0 < u \leq t\}) \geq k\}$. We are recursively adding 1 new arrival after the last arrival using the renewed intensity $\omega(\tilde{\tau}_{k-1}, \cdot)$ in (144). Define a process $\tilde{N}(t)$ by

$$(145) \quad \tilde{N}(t) = \max\{k \in \mathbb{Z}_+ \mid \tilde{\tau}_k \leq t\}, \quad t \geq 0.$$

Since $\tilde{N}(t)$ and $\tilde{\tau}_k$ are samplewise non-decreasing in t and k , respectively, $\tilde{N}(t)$ satisfies (140) and (141) with $N(t) = \tilde{N}(t)$ and $\tau_k = \tilde{\tau}_k$. $\{\tilde{\tau}_k \leq t\}$ is in

$$(146) \quad \mathcal{F}_t := \sigma[\nu(A); A \in \mathcal{B}([0, \infty)^2), A \subset [0, \infty) \times [0, t]],$$

and consequently \tilde{N} is adapted to $\{\mathcal{F}_t\}$.

In coupling the stochastic ranking process with the flow driven stochastic ranking process, we will need a representation of N by the stochastic integration with respect to ν in (143) which is,

$$(147) \quad N(t) = \int_{s \in (0, t]} \int_{\xi \in [0, \infty)} \mathbf{1}_{\xi \in [0, \omega(\tau^*(s-), s-))} \nu(d\xi ds), \quad t \geq 0,$$

where $\tau^* : \Omega \times [0, \infty) \rightarrow [0, \infty)$ is the last arrival time up to time t :

$$(148) \quad \tau^*(t) = \tau_{N(t)} = \inf\{s \geq 0 \mid N(t) = N(s)\} \in [0, t],$$

which satisfies a stochastic integration equation

$$(149) \quad \tau^*(t) = \int_{s \in (0, t]} \int_{\xi \in [0, \infty)} (s - \tau^*(s-)) \mathbf{1}_{\xi \in [0, \omega(\tau^*(s-), s-))} \nu(d\xi ds),$$

from which (147) follows.

For $t \geq t_0$ put

$$(150) \quad \Omega(t_0, t) = \int_{t_0}^t \omega(t_0, u) du.$$

We have explicit formula

$$(151) \quad \begin{aligned} \mathbb{P}[N(t) = N(s)] &= \sum_{k \geq 0} \mathbb{P}[\tau_k \leq s, t < \tau_{k+1}] \\ &= \sum_{k \geq 0} \int_{0 =: u_k < u_{k-1} < u_{k-2} < \dots < u_1 < u_0 \leq s} \\ &\quad \times e^{-\sum_{i=0}^{k-1} \Omega(u_{i+1}, u_i) - \Omega(u_0, t)} \left(\prod_{i=0}^{k-1} \omega(u_{i+1}, u_i) du_i \right), \end{aligned}$$

and for $\|\omega\| = \sup_{0 \leq s \leq t \leq T} |\omega(s, t)|$,

$$(152) \quad \begin{aligned} 0 &\leq -\frac{\partial}{\partial t} \mathbb{P}[N(t) = N(s)] \leq \|\omega\| (\mathbb{P}[N(t) = N(s)] - \mathbb{P}[N(t) = N(0)]) \leq \|\omega\|, \\ 0 &\leq \frac{\partial}{\partial s} \mathbb{P}[N(t) = N(s)] \\ &= \sum_{k \geq 0} \int_{0 =: u_k < u_{k-1} < \dots < u_1 < u_0 \leq s} w(u_0, s) \\ &\quad \times e^{-\sum_{i=0}^{k-1} \Omega(u_{i+1}, u_i) - \Omega(u_0, t)} \left(\prod_{i=0}^{k-1} \omega(u_{i+1}, u_i) du_i \right) \\ &\leq \|\omega\| \mathbb{P}[N(t) = N(s)] \leq \|\omega\|, \\ 0 &\leq s < t. \end{aligned}$$

References

- [1] H. Bauer, *Measure and integration theory*, de Gruyter Stud. Math. **26**, Walter de Gruyter & Co., Berlin, 2001.
- [2] P. Billingsley, *Convergence of probability measures, 2nd ed.*, John Wiley and Sons, New York, 1999.
- [3] P. Erdős, *On a Theorem of Hsu and Robbins*, Ann. Math. Stat. **20** (1949) 286–291.
- [4] P. Erdős, *Remark on my Paper ‘On a Theorem of Hsu and Robbins’*, Ann. Math. Stat. **21** (1950) 138.
- [5] Y. Hariya, K. Hattori, T. Hattori, Y. Nagahata, Y. Takeshima, T. Kobayashi, *Stochastic ranking process with time dependent intensities*, Tohoku Mathematical Journal **63–1** (2011) 77–111.
- [6] K. Hattori and T. Hattori, *Existence of an infinite particle limit of stochastic ranking process*, Stochastic Process. Appl. **119** (2009) 966–979.
- [7] K. Hattori and T. Hattori, *Equation of motion for incompressible mixed fluid driven by evaporation and its application to online rankings*, Funkcial. Ekvac. **52** (2009) 301–319.
- [8] K. Hattori and T. Hattori, *Sales ranks, Burgers-like equations, and least-recently-used caching*, RIMS Kokyuroku Bessatsu **B21** (2010) 149–162.
- [9] T. Hattori, *Solving the mystery of Amazon sales ranks (in Japanese)*, Kagaku Dojin, Kyoto, 2011.
- [10] T. Hattori, *Stochastic ranking process and web ranking numbers*, in *Mathematical Quantum Field Theory and Renormalization Theory*, T. Hara, T. Matsui, F. Hiroshima, eds., Math-for-Industry Lecture Note Series **30** (2011) 178–191.
- [11] T. Hattori, *Point process with last-arrival-time dependent intensity and 1-dimensional incompressible fluid system with evaporation*, Funkcialaj Ekvacioj **60** (2017) 171–212. (<http://arxiv.org/abs/1409.5117>.)
- [12] T. Hattori, *Open problems to an infinite system of quasi-linear partial differential equations with non-local terms*, Symposium on Probability Theory 2014/12/16–19, RIMS Kokyuroku **1952** (2015) 9–16.

- [13] T. Hattori, *Doubly uniform complete law of large numbers for independent point processes*, preprint (2016) <http://web.econ.keio.ac.jp/staff/hattori/LLN15r.pdf>
- [14] T. Hattori, S. Kusuoka, *Stochastic ranking process with space-time dependent intensities*, ALEA, Lat. Am. J. Probab. Math. Stat. 9(2) (2012) 571–607.
- [15] P. L. Hsu, H. Robbins, *Complete convergence and the law of large numbers*, Proc. Nat. Acad. Sci. U.S.A. **33** (1947) 25–31.
- [16] Y. Nagahata, *Tagged particle dynamics in stochastic ranking process*, Kodai Math. J. **36** (2013) 397–408.
- [17] Y. Nagahata, *Functional central limit theorem for tagged particle dynamics in stochastic ranking process*, Kodai Math. J. **36** (2013) 409–427.
- [18] M. L. Tsetlin, *Finite automata and models of simple forms of behaviour*, Russian Math. Surv. **18** (1963) 1–27.

LABORATORY OF MATHEMATICS
FACULTY OF ECONOMICS
KEIO UNIVERSITY
HIYOSHI CAMPUS
YOKOHAMA 223-8521
JAPAN

E-mail address: hattori@econ.keio.ac.jp